

Exogenous Opioids and the Human Endocrine System: An Endocrine Society Scientific Statement

Niki Karavitaki^{1,2,3}, Jeffrey J. Bettinger⁴, Nienke Biermasz⁵, Mirjam Christ-Crain^{6,7}, Monica R. Gadelha⁸, Warrick J. Inder^{9,10}, Elena Tsourdi^{11,12}, Sarah E. Wakeman¹³, Maria Zatelli¹⁴

¹Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK; ²Centre for Endocrinology, Diabetes and Metabolism, Birmingham Health Partners, Birmingham, UK; ³Department of Endocrinology, Queen Elizabeth Hospital, University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK; ⁴Pain Management and Addiction Medicine, Saratoga Hospital Medical Group, Saratoga Springs, NY, USA; ⁵Department of Internal Medicine, Division of Endocrinology, Leiden University Medical Center, Leiden, The Netherlands; ⁶Department of Endocrinology, Diabetes and Metabolism, University Hospital Basel, Basel, Switzerland; ⁷Department of Clinical Research, University of Basel, Basel, Switzerland; ⁸Endocrine Unit and Neuroendocrinology Research Center, Medical School and Hospital Universitário Clementino Fraga Filho - Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil; ⁹Department of Diabetes and Endocrinology, Princess Alexandra Hospital, Brisbane, Queensland, Australia; ¹⁰Medical School, the University of Queensland, Brisbane, Queensland, Australia; ¹¹Department of Medicine III; ¹²Center for Healthy Aging, Technische Universität Dresden, Dresden, Germany; ¹³Massachusetts General Hospital, Program for Substance Use and Addiction Service, Mass General Brigham, Harvard Medical School, Boston, USA; ¹⁴Section of Endocrinology, Geriatrics and Internal Medicine, Department of Medical Sciences, University of Ferrara, Ferrara, Italy.

Keywords: opioids, opiates, hypothalamo-pituitary, hypopituitarism, hypogonadism, bones.

Corresponding author and author to whom reprint requests should be addressed:

Niki Karavitaki, MSc PhD FRCP
Institute of Metabolism and Systems Research (ISMR)
College of Medical and Dental Sciences
University of Birmingham
IBR Tower, Level 2
Birmingham, B15 2TT, UK
Email: n.karavitaki@bham.ac.uk; Phone: +44-0121-414-3826

Abstract (max 250 words)

The use and misuse of opioids are a growing global problem. Although the effects of these drugs on the human endocrine system have been studied for decades, attention on their related clinical consequences, particularly on the hypothalamo-pituitary system and bone health, intensified over recent years. This Statement appraises research data related to the impact of opioids on the gonadal and adrenal function; whilst hypogonadism is well recognised as a side effect of opioids, the significance of their inhibitory actions on the hypothalamo-pituitary-adrenal system and the occurrence of clinically relevant adrenal insufficiency is not fully elucidated. The often-inconsistent results of studies investigating how opioids affect the secretion of growth hormone, prolactin, arginine vasopressin and oxytocin are assessed. The accumulating evidence of opioid actions on bone metabolism and their negative sequelae on bone mineral density and risk of fracture are also reviewed. In each section, available data on diagnostic and management approaches for opioid endocrine sequelae are described. This Statement highlights a plethora of gaps in research associated with the effects and clinical consequences of opioids on the endocrine system. It is anticipated that addressing these gaps will improve the care of people using or misusing opioids worldwide. The Statement is not intended to serve as a guideline or dictate treatment decisions.

Introduction

Opiates are a class of medicine derived from the opium poppy; a plant that has been used for several millennia as an analgesic by a vast number of segments of human race (1). The naturally occurring alkaloids of opium and the drugs synthesized from it are described as ‘opiates’, whilst all natural or synthetic chemicals that bind to opioid receptors are included in the term ‘opioids.’ Their first reported use was around 4000 BCE, when opium poppy (*Papaver somniferum*) was cultivated in Mesopotamia by the Sumerians, though since that time, many opioid products have been derived from opium poppy and synthetically created, with the majority produced to relieve various types of pain (1). Opioids may also be misused due to their euphoric effects. Increasing opioid use and shifts in the illicit drug supply towards high potency synthetic opioids has resulted in a public health crisis with a dramatic increase in overdose mortality (2-6).

This Scientific Statement focuses on the effects of exogenous opioids on the human endocrine system, on the evidence related to the clinical significance of these actions and on the optimal diagnostic and management approach of opioid-induced endocrinopathies. The Statement specifically depicts the landscape of opioids in cancer and non-cancer pain and of opioid use disorder (OUD), and presents data on the impact of these drugs on the hypothalamo-pituitary system and on bone health. It also identifies important gaps in our knowledge on this topic highlighting directions for future research.

Opioids in Cancer and Non-Cancer Pain and Opioid Use Disorder

Opioid Use for Analgesia

Prescription opioids are used to manage chronic or acute pain. However, the risks of prescription misuse, OUD, and overdose have been a growing concern globally (7).

US opioid prescribing catapulted in the late 1990s and into the early 2000s (2,8,9). Corresponding with the significant climb in the annual number of overall opioid prescriptions was a similarly appearing upswing of opioid-related overdose deaths (8-10). Later, as opioid prescribing rates declined, there was a transition in use and death from non-medical prescription opioid use to illicit opioids, which at first was heroin, and then, as fentanyl penetrated the drug supply, became nearly entirely supplanted by fentanyl related analogs (3,4,10-14). Three distinct waves of opioid-mortality have been described in the current overdose crisis; the first began with increased prescribing of opioid pain relievers (natural and semi-synthetic opioids and methadone) in the 1990s, the second started in 2010 and was dominated by heroin, whereas the third started in 2013 and was driven predominantly by illicitly manufactured fentanyl and related analogs (15). In recent years, co-use of stimulants with opioids has risen, leading to what some experts are now calling the fourth, polysubstance wave of the overdose crisis (12,16,17).

Because of the rise in opioid-related overdose deaths and the widespread, and often unmonitored prescribing of opioids, several guidelines and policies were developed to try to endorse monitoring strategies, optimize dosing procedures, enhance risk mitigation, and curb opioid prescribing, in an attempt to decrease overdose death rates (18-25). Specifically, the Centers for Disease Control and Prevention (CDC) published their opioid prescribing guidelines in 2016 (19), which was followed by dramatic changes in third party payer, legislative, and healthcare system level policies, often limiting overall doses of opioids that could be used for chronic non-cancer pain and emphasizing the need to utilize other modalities to treat pain (26).

According to the World Drug Report 2020, in 2018, the estimated number of people who had used opioids in the last year was 57.8 million [this figure includes those who had used opiates (30.4 million), as well as those who had misused pharmaceutical opioids] (27). Based on these data, the past-year prevalence of opioid use was 1.2% of the global population aged between 15 and 64 years; prevalence was higher than the global average in North America (3.6%), Australia and New Zealand (3.3%), Near and Middle East and South-West Asia (2.6%) and South Asia (2.0%) (27). It is of note that around 20% of the global population aged between 15 and 64 lives in South Asia and more than one third of the number of opioid users around the world live in this subregion (27). A survey of the European Pain Federation showed that in most of the 25 European countries surveyed, opioid prescriptions increased from 2004 to 2016 with a different rate between the various countries (28). In 2022, the total number of dispensed opioid prescriptions in US was 131,778,501 with an opioid dispensing rate of 39.5 per 100 persons; in 2019, these figures were 153,626,197 and 46.8 per 100 persons, respectively (10). The International Narcotics Control Board Report 2022 revealed that the highest consumption of the main opioid analgesics (codeine, dextropropoxyphene, dihydrocodeine, fentanyl, hydrocodone, hydromorphone, ketobemidone, morphine, oxycodone, pethidine, tilidine and trimeperidine) [expressed in S-DDD (Defined Daily Dose for Statistical purposes) per million inhabitants per day] is in developed countries in Europe and North America; for the year 2021, the regional averages were 17,035 S-DDD for North America, 8,721 S-DDD for Western and Central Europe and 7,146 S-DDD for Oceania, placing North America as the region with the highest consumption of opioids for pain management in the world (29).

US and the world continue to grapple with maintaining the balance between dealing with the worst overdose crisis in the history of human-kind and still considering the fact that opioids may be necessary, and therapeutically appropriate, to treat different types of pain. One of the most important ways to help ‘walk the pain management line’ is to enhance one’s understanding of opioid pharmacology and physiochemical characteristics, and how they relate to analgesic and potential adverse effects.

Opioid Use Disorder

146
147 OUD is defined as the compulsive use of opioids despite negative consequences and is diagnosed based
148 on criteria using the Diagnostic and Statistical Manual of Mental Disorders, fifth edition (DSM-5). A
149 person is diagnosed with OUD if they have experienced at least two of 11 symptoms in a 12-month
150 period and the symptoms have led to clinically significant impairment or distress (30). OUD is classified
151 as mild, moderate, or severe depending on the number of DSM-5 criteria met. OUD symptom checklists
152 have been created to support providers in making a diagnosis (31). The 11 diagnostic criteria include
153 evidence of tolerance; withdrawal; using more than intended; difficulty controlling opioid use; spending
154 more time using, recovering from, or seeking opioids; craving or a strong psychological urge to use
155 opioids; using in physically hazardous situations; using despite medical or psychiatric problems caused
156 or exacerbated by opioid use; failing to meet role obligations because of opioid use; continuing to use
157 opioids despite social problems caused by use; and giving up activities because of use (32). Tolerance
158 and withdrawal alone in the context of taking prescribed opioids does not meet criteria for OUD.

159
160 In 2021, among individuals aged 12 and older in the US, 1.1 million used heroin in the past year, 8.7
161 million reported misusing prescription opioids, 26,000 reported newly using heroin for the first time in
162 the past year, and 5.6 million people met criteria for OUD (33). Among people who reported misusing
163 prescription opioids, 64% either used more of their own prescription or used opioid pain relievers
164 without a prescription for the purpose of relieving physical pain. However, 11% used them to feel good
165 or get high and 7% to relax (33). Globally, in 2016 there were an estimated 26.8 million people alive
166 with OUD (34). In 2017 alone, the estimated cost in the US related to OUD and fatal opioid overdose
167 was one trillion dollars due to reduced quality of life and the value of life lost from fatal overdose (35).

168
169 Despite touchpoints with the healthcare system most people with OUD do not receive treatment and
170 racial and ethnic disparities persist in treatment access (36-41). Racial and ethnic disparities are also
171 widening in the overdose crisis, with the greatest rates of increase in opioid-related deaths between
172 2019 and 2020 in US seen among Black Americans and American Indian/Alaskan Native populations
173 (absolute rate change of drug overdose deaths between 2019 and 2020: Non-Hispanic Black persons

11.9, American Indian/Alaskan Native populations 10.2, White persons 5.5) (42). Social determinants of health exacerbate these racial and ethnic disparities, with greater racial and ethnic disparities in overdose seen in communities with the highest income inequality (42).

OUD is a treatable health condition for which effective treatments exist. Long-term medication treatment with methadone (a full opioid agonist) and buprenorphine (a partial opioid agonist) has been shown to improve remission rates, quality of life and overall health and to decrease ongoing opioid use, overdose, and mortality (43,44). In addition to methadone and buprenorphine, a third Food and Drug Administration (FDA)-approved medication for OUD is extended-release naltrexone, an opioid antagonist, which is a second line treatment option for the prevention of recurrence of opioid use among individuals who have completed opioid withdrawal management.

Pharmacology of Opioid Analgesics

Opioids primarily exert their physiologic and pharmacologic effects by modulating the endogenous opioid receptor system. This system consists of four major opioid receptors (G protein-coupled) including mu-opioid receptors (MORs), kappa-opioid receptors (KORs), delta-opioid receptors (DORs), and opioid-like receptor one (also known as nociceptin) (44-46). In addition to these main groups, there are several subtypes of each receptor, and the entire receptor system can be susceptible to pharmacogenomic abnormalities, all of which may impact an opioid's ability to modulate such receptors and can help explain individual and site variabilities regarding specific responses to opioids and their overall effects (44-46). While all four receptor types play important roles from a physiologic perspective, most data (especially from an endocrine perspective) continue to revolve around opioids impact of MORs and the response induced by that.

Most opioids used today act by fully agonizing MORs. All full opioid receptor agonists activate opioid receptors, thereby inducing a conformational change with the receptor and causing several intracellular events that inevitably lead to physiologic effect (47,48). Other opioids are considered ‘partial agonists’

at MORs, including buprenorphine and butorphanol, whereby binding to MORs still activates those receptors, albeit to a lesser degree than full agonists (47,48). The general ‘amount’ by which receptors are activated by an individual opioid (or any compound) is considered its intrinsic activity (or biological stimulus), and it is usually measured in *in vitro* studies (47,48). It is important to note that these studies do not always completely correlate with efficacy/clinical response which revolves around a particular endpoint (47,48). Finally, there are opioids that antagonize MORs, such as naloxone, naltrexone, methylnaltrexone, whereby binding to MORs simply blocks further activity and does not yield any conformational receptor change nor response (47,48) (Table 1).

Table 1: Properties of Commonly Used Opioid Agonists. From Ref (49).

Opioid	Binding affinity toward MOR	Relative lipophilicity ^a	MME dose ^b	Approximate equivalent dose (mg) ^c
Sufentanil	+++++	+++++	---	0.02
Buprenorphine	+++++	+++++	1.8 (transdermal patch, µg/h)	0.3
Hydromorphone	++++	++	4.0	1.5
Oxymorphone	++++	+	3.0	1.5
Levorphanol	++++	++++	11.0	2–3
Morphine	+++	+	1.0	10
Fentanyl	+++	+++++	2.4 (transdermal patch, µg/h)	0.1
Oxycodone	++	+	1.5	4.5
Codeine	+	++	0.15	30–60

This table should not be used to convert one opioid to another for clinical use.

MME morphine milligram equivalent; MOR mu opioid receptor.

^a Based on log P, which corresponds to the logarithm of the ratio of the concentrations of the studied compound in octanol and in water: $\log P = \log (C_{oct}/C_{water})$. (Values obtained from reference [(50)]).

^b From the Centers for Disease Control and Prevention <https://cdc.gov/drugoverdose/training/dosing/accessible/index.html> (51).

Morphine Milligram Equivalent (MME) Calculators:

<https://www.mdcalc.com/morphine-milligram-equivalents-mme-calculator>

<https://agencymeddirectors.wa.gov/Calculator/DoseCalculator>

^c (52).

Opioid receptors are expressed throughout the body, including the central nervous system (CNS), peripheral nervous system and enteric nervous system, amongst other sites. Opioid actions and clinical responses generally depend on where and which receptor the opioid is activating. Regarding pain pathways, opioid receptors are expressed throughout the descending pain pathways in the cortical and

midbrain structures, periaqueductal gray area, limbic system, medulla locus coeruleus dorsal root ganglion, and throughout peripheral afferent neurons as well (45,46,53). Binding to MORs throughout those areas is what is primarily associated with pain attenuation and analgesia.

Aside from MORs presence throughout various pain pathways, they are also present throughout other regions of the body, and their location may help explain specific opioid non-analgesic physiologic effects. Some of the most common opioid-related side effects include delayed gastric emptying, slowed gastro-intestinal transit, and constipation, which are all caused by MOR activation throughout the gastrointestinal tract leading to reduction in intestinal secretions, greater fluid efflux from the gastric lumen, anal sphincter paralysis, and reduced peristalsis (45,46). Opioids can also be associated with nausea and vomiting because of their activation of the medullary chemoreceptor trigger zone within the CNS (45,46). Opioid activity throughout the ventral tegmental and nigrostriatal cortex, and corresponding communication with other sections of the CNS like the amygdala, are some of the mechanisms thought to allow for opioid-induced euphoria and may be involved in the development of addiction (45,46,54). Perhaps the most dangerous side effect of opioid use involves their activity within the pre-Botzinger complex and parabrachial nucleus, as this is what is associated with suppression of respiratory function and attenuation of the central response to rising carbon dioxide levels, leading to potential cardio-pulmonary collapse and death (45,46,54). Opioids can also have actions on the hypothalamic-pituitary system, which are one of its most underappreciated side effects and are covered in this Statement.






Distinguishing Between Opioids


There are several key differences between different opioid compounds that are important to recognize, especially as they relate to varying clinical effects and overall potencies. First, it is essential to understand that exogenous opioids are generally classified by how they are created, and then by their base chemical structure which helps determine almost everything about the drug itself. Natural opiates are considered alkaloids that are derived directly from the opium poppy itself including morphine,

codeine, and thebaine (1). Semi-synthetic opioids are opioids that are synthesized directly from natural opiates, and include several common ones, such as hydrocodone, oxycodone, hydromorphone, buprenorphine, and heroin (diacetyl morphine), amongst others (1). Fully synthetic opioids are opioid compounds that are synthesized completely in the lab, such as methadone, fentanyl, tapentadol, and tramadol, amongst others (1). All these opioids can then be further categorized based on their base chemical structures, of which there are five major groupings. These include the phenanthrenes, benzomorphans, phenylpiperidines, diphenylheptanes, and phenylpropyl amines (Table 2).

Table 2. Chemical classes of opioids.

Chemical Classes of Opioids (Updated 11/1/2020)

PHENANTHRENES	BENZOMORPHANS	PHENYLPIPERIDINES	DIPHENYLHEPTANES	PHENYLPROPYL AMINES
				
MORPHINE	PENTAZOCINE	FENTANYL	METHADONE	TRAMADOL
Buprenorphine* Butorphanol* Codeine Dextromethorphan* Dihydrocodeine Heroin (diacetyl-morphine) Hydrocodone* Hydromorphone* Levorphanol* Methylnaltrexone** Morphine (Opium, conc) Nalbuphine* Naloxone* Naloxegol* Naltrexone** Oxycodone* Oxymorphone*	Pentazocine	Alfentanil Fentanyl Remifentanyl Sufentanil Meperidine Diphenoxylate ^a Loperamide ^a	Methadone Propoxyphene	Tapentadol Tramadol
		Fentalogues Illicit Fentanyl Analogues		
		Furanyl fentanyl Acetyl fentanyl Fluoro-fentanyl Carfentanil Others ^b		
CROSS-SENSITIVITY RISK				
PROBABLE	POSSIBLE	LOW RISK	LOW RISK	LOW RISK
*Agents lacking the 6-OH group of morphine, possibly decreases cross-tolerability within the phenanthrene group				
**6-position is substituted with a ketone group and tolerability is similar to hydroxylation				



Courtesy of Jeffrey Bettinger, all rights reserved.

^a Previously incorrectly listed as "Benzomorphans".

^b (55).

While an in-depth comparison between each individual opioid is outside the scope of this Statement, it is important to review some of the most pertinent differences. Certainly, opioid pharmacokinetic and pharmacodynamic variability is mainly driven by differences in their physiochemical characteristics. Considering most of the opioid-induced analgesia occurs because of opioid actions throughout the CNS

and peripheral nervous system, thus requiring transport through the blood brain barrier (BBB), opioids with more lipophilic chemical structures more easily transport through the BBB to their sites of action (56-58). This is why lipophilicity is one of the most critical characteristics when assessing measures of potential potency around clinical endpoints. Such opioids with greater degrees of lipophilicity include fentanyl, sufentanil, buprenorphine, and hydromorphone (56-58) (see Table 1 physiochemical properties). In addition to lipophilicity, another important physiochemical characteristic is an opioid's binding affinity, which is defined as the thermodynamically driven chemical attraction between a substance and a receptor, and usually represented by its K_i value (59). Opioids with lower K_i values have greater degrees of binding affinity toward opioid receptors. Once again, buprenorphine, sufentanil, hydromorphone, and oxymorphone have some of the greatest binding affinities toward MORs (59). Not only can these characteristics help define doses used for different opioids from an analgesic perspective (as well as formulations required), but they can help delineate between analgesic efficacy and toxicity risk; including from an endocrine perspective. Buprenorphine, for example, is a lipophilic opioid with significant binding affinity toward MORs and is considered to be highly potent, as low doses (usually in micrograms) are required to produce cellular responses (48,57,60). However, because it is a partial agonist (has lower intrinsic activity at MORs), it does not produce as much of a clinical response as full MOR agonists, particularly from a toxicity perspective (48,57,60). Thus, comparatively, buprenorphine is associated with plateaued respiratory depressive effects and results in less gonadal axis suppression. Finally, half-life is another key differentiator when it comes to opioid-induced endocrine effects.

Hypothalamo-Pituitary-Gonadal axis

Effects of opioids

The inhibitory actions of opioids on the hypothalamo-pituitary-gonadal axis and the subsequent resulting hypogonadism are well recognised. Opioids inhibit pulsatile gonadotropin-releasing hormone (GnRH) secretion from the hypothalamus and the secretion of gonadotropins [mainly luteinising hormone (LH) and to a lesser extent follicle-stimulating hormone (FSH)] from the pituitary gland and of sex steroids from the gonads (61-64). They also can inhibit the secretion of dopamine from the hypothalamus leading to hyperprolactinemia, which further contributes to the suppressive action on the hypothalamo-pituitary-gonadal axis (65). Direct negative effects of opioids on testicular function and sperm quality have also been proposed (66-68).

Prevalence of hypogonadism in males on opioids has been reported within variable rates, depending on type, dose, route, and duration of administration of the opioid, diagnostic approach for the confirmation of hypogonadism, as well as on the patient group assessed (65,69). The first studies on the negative effects of opioids on the hypothalamo-pituitary-gonadal axis involved mainly people addicted to heroin and patients on methadone maintenance treatment, with hypogonadism reported in 40-85% of males (70). In female heroin addicts, amenorrhea and galactorrhea have also been described, although this effect has not been systematically looked at in this group (71). In a meta-analysis of 15 studies with a total of 3250 patients (99.5% males) receiving chronic various opioids, the weighted mean rate of hypogonadism (based on a single morning or random testosterone measurement) was 63% [95% Confidence Interval (CI): 55-70%] (72). In another systematic review including 165 women aged 18-55 years on oral or intrathecal opioids for chronic non-cancer pain, 23-81% of them had oligo/amenorrhea (69). In a case control study of 21 women on sustained action oral or transdermal opioids for non-malignant pain and 16 women not on opioids (but not matched for confounding factors) aged between 30 and 50 years, the levels of estradiol, testosterone and dehydroepiandrosterone (DHEAS) were significantly lower in opioid users (73). Furthermore, reduced levels of gonadotropins

have been found in post-menopausal women on intrathecal opioids for non-malignant pain (74). It should be noted that when interpreting data on the prevalence of opioid-induced hypogonadism in patients with chronic non-malignant pain, other confounding factors, like effects of pain, other comorbidities (e.g., obesity), and/or concurrent medications need to be taken into account. Hypogonadism has also been reported in patients receiving opioids for cancer-related pain (75,76) but in this group, in addition to the previously mentioned factors, the underlying malignant disease, chemotherapy, anorexia/cachexia and psychological stress may have a further negative impact on the axis (77). In a study of male cancer survivors (disease-free for more than one year) using opioids [morphine milligram equivalent (MME) daily dose ≥ 200 mg], hypogonadism was found in 90% of them (18/20), as compared to 40% (8/20) of subjects in a matched control group. Median testosterone was significantly lower in the opioid group, although testosterone measurements were not performed on early morning samples, potentially affecting the reliability of the hypogonadism diagnosis (78).

The inhibitory actions of opioids on the hypothalamo-pituitary-gonadal axis occur as soon as they are administered (74,79,80) and cessation or reduction in their dose leads to recovery of the axis (79-82), the time course of which is variable (80,83-85). The severity of hypogonadism depends on the dose and type of opioid (86-88). Long-acting opioids are more likely to cause androgen deficiency than the short acting ones (89), and transdermal fentanyl, oral methadone, and oral oxycodone have been reported to have higher odds of androgen deficiency compared to hydrocodone (90). Buprenorphine, a partial opioid receptor agonist, is associated with milder effects on the hypothalamo-pituitary-gonadal system (91,92).

Clinical significance

Typical clinical manifestations of gonadal hormone deficiency have been described in patients with opioid-induced hypogonadism (erectile dysfunction, decreased muscle mass in men, oligo/amenorrhea in women, decreased libido, infertility, bone loss, depression in both sexes) (65,69,72). Data on the impact on fertility are limited (93) and adverse effects of these drugs on semen quality have been

suggested (68). Hypogonadism may remain undiagnosed, particularly in men, possibly due to lack of reporting of the relevant symptoms by the patients and to underappreciation of this side effect by the health care professionals.

Interestingly, it has been proposed that opioid-induced androgen deficiency might impair pain control and lead to hyperalgesia (94) .

Diagnostic approach

Timely diagnosis of hypogonadism is of major importance. In accord with the Endocrine Society guidelines, screening should be avoided in patients on short term opioid regimens (95). In contrast, patients on long-term opioid treatment (suggested interval longer than 3 months) need to be informed about the risk of hypogonadism and encouraged to report relevant symptoms. Also, manifestations of hypogonadism need to be enquired by clinicians and if high index of suspicion, diagnostic investigations should be performed as per current guidelines (96).

Management

Given that discontinuation or reduction of the opioid dose reverses their inhibitory effects on the hypothalamo-pituitary-gonadal axis (79-82), this approach is considered to be the first management step, when possible (in parallel, further management options for the pain should be discussed and provided by the relevant experts). The time course of the recovery has not been clearly established and the impact of opioid tapering on hypogonadism has not been explored. Change to buprenorphine, an opioid with milder or no suppressive effect on the axis, could also be another approach. If these are not feasible, gonadal hormone replacement therapy needs to be considered aiming to avoid the negative sequelae of untreated hypogonadism (70,96).

The effects of gonadal hormone replacement in patients with opioid-induced hypogonadism have been investigated in a small number of studies involving only males and have provided some promising outcomes. Nonetheless, most are of rather short duration posing challenges in the assessment of the safety of this treatment. An open-label pilot study of 23 men on oral sustained release oxycodone or oral methadone or transdermal fentanyl or oral sustained release morphine sulfate for pain and with morning free testosterone <50 pg/ml (normal range 52-280 pg/ml) examined the effects of testosterone patches administered for 24 weeks (5 mg daily for the first 12 weeks and 7.5 mg daily for the second 12 weeks); in the 16 patients who finally completed the trial, improvement in their androgen deficiency symptoms, sexual function, mood, depression, and hematocrit was demonstrated (97). In a study of 9 males on epidural morphine for non-cancer chronic pain and 8-9 am testosterone less than 2-3 ng/ml (in at least two determinations in the previous 3-4 months, normal range 3.5-8.5 ng/ml), a sachet of testosterone gel (containing 50 mg testosterone in 5 g gel) was administered daily for 12 months; this treatment was associated with improvement in the sexual dimension of the Aging Males' Symptoms scale and in the mental index of the Short-Form (SF-36) Health Survey, but not in the Profile of Mood State subscale scores or Centre for Epidemiological Studies Depression Scale ratings (98).

In a randomized, double-blind, placebo-controlled parallel-group trial of 65 men on at least 20 mg daily of hydrocodone (or MME dose of other opioid) during a minimum of four weeks for chronic non-cancer pain and with morning serum total testosterone below 350 ng/dl (measured by liquid chromatography–tandem mass spectrometry) were randomly assigned to receive either 5 g of a transdermal testosterone gel or placebo gel, applied once daily for 3 months. Two weeks after randomization, if total testosterone was below 500 ng/dl, the dose of the testosterone gel was increased to 7.5 g daily. Those on transdermal testosterone showed greater improvement in sexual desire, fat mass, as well as in pressure pain threshold at the thumb and mechanical hyperalgesia compared with the patients on placebo (99). In a double-blind, placebo-controlled study, 41 men with non-malignant pain on opioids for at least three months at a minimum MME dose of 50 mg daily and with 8-10 am total testosterone below 12 nmol/L (346 ng/dL) were randomized to have intramuscular testosterone undecanoate 1000 mg or placebo injection at baseline and at 6 and at 18 weeks; at 6 months, in those treated with testosterone, increased lean body

mass, and reduced total fat mass, but no change in pain perception were observed (100). In a study of 27 males on opioids for chronic non-cancer pain and opioid-induced hypogonadism (total testosterone below 300 ng/dL), 11 of them were offered testosterone supplemental therapy (topical or injectable or subcutaneous implant with no further details on this treatment) and their outcomes were compared with those 16 untreated (101). Data were collected at baseline and follow-up at 6 monthly periods over 18 months. The testosterone treated group showed significant improvement in the Numerical Rating Scale pain score and a reduction in the opioid requirements, as quantified by daily MME dose (mean daily MME dose decreased by 21 mg in this group and increased by 2.5 mg in the non-treated one, $p<0.05$), and also an improvement in hypogonadal symptoms [assessed by the Androgen Deficiency in Aging Males (ADAM) and the International Index of Erectile Function (IIEF-5) questionnaires] (101). In a double-blind, placebo-controlled study involving men with non-cancer pain treated with opioids for at least three months on a MME dose of at least 50 mg daily and with total testosterone below 12 nmol/L (measured two times between 8 and 10 am in the fasting state by liquid chromatography-tandem mass spectrometry), 20 patients were randomly assigned to receive testosterone undecanoate injection 1000 mg and 21 participants to receive placebo (102). The injections were offered at the time of randomization, at 6 and at 18 weeks, and muscle function, gait performance and body composition were measured prior to and after 24 weeks of intervention. The trial was completed by 38 participants and 6 of them were not able to complete all tests of muscle function. The testosterone treatment group showed significant increase in the lean body mass, leg lean mass, body weight and a reduction in the total fat mass compared to placebo. Despite these changes, there was no difference in the muscle function or gait performance. The authors suggested that people with chronic pain are more likely to exhibit sedentary behavior and, therefore, the absence of exercise/high intensity physical activity may explain the lack of association in the observed changes in body composition and muscle function in the testosterone treated participants (102). Finally, in a placebo-controlled, double-blind randomized study, 64 non-diabetic males aged between 18 and 64 years on opioid analgesics for chronic non-cancer pain (at least 20 mg hydrocodone or MME dose of another opioid) and morning total testosterone below 12 nmol/L (measured by liquid chromatography-tandem mass spectrometry) were randomised to 14 weeks of transdermal testosterone gel (5 gr) or placebo gel daily (two weeks after randomization, if the serum

total testosterone level was below 17 nmol/l, in which case, the dose of testosterone gel was increased to 7.5 g daily); at the end of the study period, changes in lipid profile, fasting glucose and insulin, homeostatic model assessment for insulin resistance and C-reactive protein were similar from baseline to the end of treatment in both arms of the study (103). Overall, treating hypogonadism with testosterone in men leads to improvement in androgen deficiency symptoms including sexual function, to reduction in fat mass and increase in lean body mass.

Hypothalamo-Pituitary-Adrenal Axis

Effects of opioids

Acute intravenous administration of endogenous opioids, such as beta-endorphin (104), enkephalin analogues (105) and pharmacological opioids, such as morphine (106), demonstrates an acute inhibitory effect on adrenocorticotrophic hormone (ACTH) and cortisol levels in humans.

Several early studies examined the hypothalamo-pituitary-adrenal axis of heroin and methadone users, with some variable results (107-109). In a study of 8 pain patients published in 1997, up to 12 weeks of oral sustained release morphine at daily doses ranging from 30-240 mg suppressed basal ACTH and cortisol levels; in two participants who had 100 mcg of human corticotropin-releasing hormone (CRH) injected (which is not a standard test for adrenal insufficiency), ACTH and cortisol concentrations showed sufficient increase (110). A landmark study published three years later, investigating the effects of intrathecal opioids using the insulin tolerance test (ITT) (the gold standard for the diagnosis of secondary adrenal insufficiency), showed that 15% (9/61) of the patients on intrathecal opioids for non-malignant pain had secondary adrenal insufficiency (74). Later, cases of symptomatic adrenal insufficiency in people taking opioids began to be described (111-113). However, a 2009 review noted that the clinical significance of the inhibitory effect of opioids on cortisol was unclear and that the clinical consequences of the low cortisol had not been established (114). Soon after, studies on larger cohorts looking at the prevalence of secondary adrenal insufficiency in people using oral or transdermal opioids for chronic non-malignant pain began to be reported (115-118). These studies were characterized by a high degree of variability in how opioid-induced secondary adrenal insufficiency was defined, and in one of them using the CRH test, in fact, a higher prevalence of secondary adrenal insufficiency in controls with chronic pain not treated with opioids was described (118). A systematic review and meta-analysis examined 52 studies assessing the endocrine effects of opioids (72). The authors included only five reports investigating hypocortisolism in the meta-analysis. These studies were heterogenous, with different studies examining cancer and non-malignant pain patients, different

methods of assessing the hypothalamo-pituitary-adrenal axis while two included intra-thecal opioid-treated patients. Among these reports, there was a weighted mean percentage of 15% [95%CI: 6-28%] with secondary adrenal insufficiency (72).

An opioid dose-dependent effect on the diagnosis of opioid-induced secondary adrenal insufficiency has been suggested. In a study assessing the effects of long-term oral or transdermal opioids in patients with non-cancer pain, an overall rate of secondary adrenal insufficiency of 22.5% (9 out of 40) was reported (defined as failing either an ACTH1-24 stimulation test or an overnight metyrapone test) (116). Using the ACTH1-24 stimulation test alone, the rate of secondary adrenal insufficiency was 15%. No patient on a MME dose of below 60 mg per day showed a suboptimal response (116). In another retrospective cohort study comprising of 40 patients considered to have opioid-induced secondary adrenal insufficiency, the minimum MME dose was also 60 mg daily (119). In addition, all cases of patients on long term oral and/or transdermal opioids for chronic pain found with blunted response to the ACTH1-24 test (3 out of 48, 6%) were on a dose higher than the MME of 60 mg per day (115). However, other studies have reported secondary adrenal insufficiency in people on lower opioid doses, including a cross sectional one examining the hypothalamo-pituitary-adrenal axis function of 102 patients on various opioids (117). Nine patients (9%) were diagnosed with secondary adrenal insufficiency [6 based on a combination of clinical assessment, basal morning cortisol, ACTH and DHEAS levels and three after an ACTH1-24 stimulation test], and their minimum MME dose was 20 mg daily. Overall, there was a significant difference in cumulative opioid exposure but not daily dose, between patients considered to have secondary adrenal insufficiency *versus* those with a normal hypothalamo-pituitary-adrenal axis (117).

The finding of a possible but inconsistent dose-dependent effect is compatible with the pharmacodynamics, pharmacokinetics, and pharmacogenomics of opioids. In a study of cancer patients, there was a two-fold difference in the steady state oxycodone concentration amongst 10 patients taking an oral identical dose (120). In another study examining pharmacogenomics, it was shown that inherited polymorphisms in the CYP2D6 enzyme determined opioid metabolism (121). In this study, 54% of

patients were extensive metabolizers, 41% intermediate and 5% poor metabolizers. Poor metabolizers had the highest steady state drug concentrations. There was a clear relationship between steady state drug concentration and degree of analgesia, and 80% of those reporting side effects had impaired CYP2D6 metabolism (121). Extrapolating these studies to the opioid effects on the hypothalamo-pituitary-adrenal axis, it is a reasonable hypothesis that a concentration threshold may exist for hormonal inhibitory effects. In turn, the concentration depends on several factors, such as drug dose, absorption, and metabolism. This potentially explains some of the variability in dose threshold which has been reported to be associated with opioid-induced secondary adrenal insufficiency. The variable prevalence of opioid-induced secondary adrenal insufficiency may in part relate to the number of poor metabolizers in the generally small cohorts. The key point is that there is unlikely to be a simple relationship between drug dose, plasma drug concentration and hypothalamo-pituitary-adrenal axis inhibition across a broader cohort of patients due to individual differences in absorption and metabolism. The interaction between drug concentration and opioid receptor expression then finally determines the extent of hormonal suppression.

Clinical significance

Despite the millions of people worldwide using oral and transdermal opioids, the number of those diagnosed with opioid-induced secondary adrenal insufficiency is low, and there have been only very few published cases of adrenal crisis (112,119,122). As well as common symptoms of cortisol deficiency, such as fatigue, nausea/vomiting, musculoskeletal pain, weight loss, headache, and lightheadedness (119), less common but well recognised manifestations, such as hypercalcemia (123) and hypoglycemia (124) have been described. Both opioid use and the pain syndromes may result in a broad spectrum of symptoms and signs which impact health-related quality of life, including physical function, sexual function, mental health, and fatigue hindering the interpretation of the relationship between health-related quality of life and hormonal data (116,124). Nevertheless, there is evidence that secondary adrenal insufficiency (in the context of pituitary disease) itself impacts aspects of health-related quality of life, including pain, which may be alleviated by increased glucocorticoid doses (125).

Diagnostic approach

In routine clinical practice, the most frequently utilised tests of hypothalamo-pituitary-adrenal axis function are the basal morning cortisol and the ACTH1-24 stimulation test (126). Low serum DHEAS has also been proposed as an adjunct test for adrenal insufficiency (127). As previously mentioned, ITT is considered the gold standard for diagnosing secondary adrenal insufficiency, although it has some limitations (128). There are data indicating that the overnight metyrapone test is a more sensitive test for secondary adrenal insufficiency than the ACTH1-24 stimulation test (129). However, a prospective study comparing the diagnostic accuracy of the ACTH1-24 stimulation test, overnight metyrapone test and ITT to diagnose secondary adrenal insufficiency after pituitary surgery found no significant differences between them, when requirement for glucocorticoid replacement at 6 months was the critical end point (130).

Most of the published studies examining the prevalence of secondary adrenal insufficiency amongst opioid users have utilised the measurement of morning cortisol with or without the 250 mcg ACTH1-24 stimulation test. Lamprecht *et al.* used morning cortisol and the 250 mcg ACTH1-24 stimulation test (applying a locally validated cut off for the 60-minute cortisol), as well as the overnight metyrapone test, although due to the reluctance of the patient group to attend morning appointments, not every participant with a morning cortisol of below 9.1 mcg/dL (250 nmol/L) had both dynamic tests (116). The cut-off for a “normal” morning cortisol varied with different studies and cortisol assays from 3 mcg/dL (83 nmol/L) (131) to 10 mcg/dL (276 nmol/L) (117). The two studies by Li *et al.* used a combination of morning cortisol, ACTH and DHEAS, suggestive symptoms and an ACTH1-24 stimulation test in selected cases, and their sampling window was between 7am and 10am (117,119). There is a considerable diurnal decrease in cortisol concentrations across this time frame. In a study where cortisol sampling was undertaken every 10 minutes for 2h between 7am and 9am, there was a median difference of 6.4 mcg/dL (177 nmol/L) between an individual’s highest and lowest cortisol across the two-hour window (130). Thus, there is a significant physiological disparity between a

“morning” cortisol drawn at 7.30am compared to 9.30am. The heterogeneity in the diagnostic criteria for opioid-induced secondary adrenal insufficiency, cortisol assay variability and a wide sampling window make direct comparison of morning cortisol levels between studies difficult. For example, in a study which included people both on and not on opioids, a cortisol of below 10 mcg/dL (276 nmol/L) was 100% sensitive to diagnose adrenal insufficiency in outpatients, but the specificity was only 54% (131).

There are no universally agreed upon criteria for the diagnosis of opioid-induced adrenal insufficiency. Manifestations of secondary adrenal insufficiency include fatigue, nausea, vomiting, postural hypotension/light-headedness, cognitive decline, muscle weakness and arthralgia/myalgia, but these may also be observed in patients on opioids with normal adrenal function. As per Endocrine Society’s guideline, the suggested approach would be to screen with a morning cortisol between 8am and 9am (96); if this is less than a pre-defined threshold consistent with the local cortisol assay, to proceed with an ACTH1-24 stimulation test, again using an assay-specific cut-off for the 30- or 60-minute cortisol (131) and keeping in mind its limitations for the diagnosis of recent-onset adrenal insufficiency. The frequency of the hypothalamo-pituitary-adrenal axis assessment in patients on opioids is not known but occurrence of manifestations of cortisol deficiency should prompt investigations for the integrity of the cortisol reserve. It is important to consider that a patient may have other causes of secondary adrenal insufficiency, such as structural pituitary disease, pharmacological doses of glucocorticoids for a variety of medical conditions or immune checkpoint inhibitors.

Management

Studies reporting on outcomes of patients with opioid-induced secondary adrenal insufficiency following glucocorticoid replacement are scarce. Li *et al.* commenced 95% of patients who met their diagnostic criteria for opioid-induced adrenal insufficiency on hydrocortisone and the remainder on prednisone (119). They reported some improvement in possible manifestations of adrenal insufficiency in 70% of those evaluated, however, analysing the individual symptoms, none demonstrated more than 50% of patients improving with glucocorticoids. The closest was fatigue, where 14 out of 29 patients

610 reported subjective improvement. Relevant but non-specific symptoms, such as headache, abdominal
611 pain, nausea, vomiting, and light-headedness were improved in less than 25% of those presenting with
612 them (119). As this was an uncontrolled study and these symptoms/signs are common in patients on
613 opioids with normal adrenal function, it is not possible to definitively attribute any clinical improvement
614 to the glucocorticoid replacement. There is only a single randomized double-blind cross-over trial of
615 glucocorticoid replacement in opioid-induced secondary adrenal insufficiency which utilized a reduced
616 cortisol response to a cold pain stimulus as the inclusion criterion (this test is not validated for the
617 diagnosis of adrenal insufficiency) (132). Five out of 17 participants had a morning cortisol below 5
618 mcg/dL (138 nmol/L), but 9 out of 10 in the trial who underwent a 1 mcg ACTH1-24 stimulation test
619 had a normal cortisol response. Therefore, while the participants had low morning cortisol and a reduced
620 hypothalamo-pituitary-adrenal axis response to a cold pain stimulus, most did not have adrenal
621 insufficiency as defined by the ACTH1-24 stimulation test. It is possible the effect of the opioid itself
622 diminishes the painful stimulus resulting in a reduced endogenous axis response. Nevertheless, the
623 study found that physiological glucocorticoid substitution had benefits in improving vitality and
624 perceived pain compared to placebo (132).

625
626 Normalization of hypothalamo-pituitary-adrenal axis function can occur rapidly following opioid dose
627 reduction. Unlike the classical scenario of glucocorticoid-induced axis suppression which can take
628 months or even years to fully recover, normal ACTH and cortisol parameters may occur within days to
629 weeks of reducing opioid doses (112,124). In a group of 7 patients, 6 of them successfully weaned off
630 opioids completely and were able to cease their hydrocortisone (117). Therefore, the primary action for
631 managing opioid-induced secondary adrenal insufficiency should be an attempt at opioid withdrawal.

632
633 There are no data to underpin a consensus on the need for daily physiological glucocorticoid
634 replacement but for those who are unable to discontinue their opioids or wean to a dose below the
635 threshold which results in opioid-induced secondary adrenal insufficiency, physiological glucocorticoid
636 replacement may be considered. In these cases, the response to treatment needs to be monitored to
637 identify if this had led to clinical improvement. The hypothalamo-pituitary-adrenal axis should be re-

evaluated after reduction or discontinuation of the opioid. There is a paradox of the prevalence of opioid use, often at high dose, yet a relatively low percentage of people with biochemically diagnosed secondary adrenal insufficiency and only very rare instances of adrenal crisis. One possible explanation is that opioids may largely affect basal ACTH/cortisol secretion but less commonly impede the physiological response to stress; therefore, the user is largely protected from the usually very serious consequences of adrenal insufficiency. Consequently, one could consider as a safe and reasonable approach in selected cases to prescribe stress dosing for times of illness but not daily maintenance dosing. Patient education on sick day rules and stress steroid doses should follow current guidelines (96). However, standard management principles applied to pathological primary and secondary adrenal insufficiency may not be universally inferred to opioid-induced secondary adrenal insufficiency. Demonstration of benefit through an appropriately powered double-blind randomized controlled trial of maintenance hydrocortisone *versus* placebo, with provision for standard stress dosing in the placebo group in the event of significant stress or illness, would answer this question.

Growth hormone axis

Effects of opioids

Studies evaluating the effects of opioids on the growth hormone (GH) axis in humans are limited and data supporting a robust and consistent effect are insufficient (70).

Single dose administration of opioids has a dose-dependent stimulatory action on the GH axis (133). This acute effect seems to be mediated through the GH-releasing hormone (GHRH) (134). Other mechanisms may also be implicated, as when a Met-enkephalin analogue G-DAMME was administered in combination with a maximally stimulatory dose of a GHRH analogue in healthy men, an enhancing effect of the GHRH-induced GH release was observed (135).

In contrast, data on the long-term impact of opioids on the somatotroph axis are contradictory. In a systematic review, the effects of opioids on the GH axis were reviewed from four studies involving 225 patients with either cancer or non-cancer pain (72,74,75,118,136). The limited number of included reports and the different protocols used to assess GH deficiency did not allow definite conclusions to be obtained (72). The study with the largest number of cases looked at 72 non-cancer pain patients who received intrathecal opioids (morphine or hydromorphone) and 20 controls (patients with non-cancer pain of similar intensity with the opioid group but not treated with these agents) (74). The opioid treated group had lower mean serum insulin like growth factor type I (IGF-I) ($138.5 \text{ mcg/L} \pm 64.1$) in comparison to the controls ($162.0 \text{ mcg/L} \pm 55.3$; $p=0.045$), as well as lower IGF-I standard deviation (SD) score (-0.53 ± 1.45 and 0.57 ± 1.00 ; $p=0.002$). Twelve opioid-treated patients (17%) had an IGF-I SD score below -2 SD, as opposed to no such case in the control group. The peak GH during the ITT was significantly lower ($p=0.010$) in the opioid group ($n=62$; $14.5 \text{ mcg/L} \pm 12.7$) compared to controls ($n=18$; $20.9 \text{ mcg/L} \pm 11.5$). Nine opioid-treated patients (15%) had a peak GH value consistent with severe GH deficiency (below 3 mcg/L), a finding not demonstrated in any control patient. These results

suggest that up to 17% of the patients treated with intrathecal opioids may develop GH deficiency (74). In a study of 40 patients on oral or transdermal opioids for chronic pain (for more than 6 months at a MME dose ≥ 25 mg) and 25 controls not taking opioids, the opioid users had lower IGF-I [138 mcg/L (99-161) *versus* 168 mcg/L (138-214); $p=0.004$], although none of them had levels below the lower limit of the reference ranges (116). Another study evaluated 39 non-cancer pain patients who received different oral opioids and 20 controls (118); GH stimulation test was not performed but mean serum IGF-I was lower in the opioid-treated group, although statistical significance was not attained. In a report comparing the effects of intrathecal and oral morphine, 18 intrathecal opioids users, 18 oral opioids users and 18 controls were evaluated (136). Serum IGF-I range was not different between the groups. Serum IGF-I levels below -2 SD were detected in 44% of the oral opioid users, in contrast to 28% of the intrathecal opioid users and controls. However, this difference was not statistically significant. Mean peak GH during ITT did not differ between the groups and, although also not statistically significant, peak GH below 3.2 mcg/L was detected in 11% of the intrathecal opioid users (136). In a study of 19 patients with cancer pain treated with several opioids, only measurements of basal serum GH levels were provided; given that this is not the appropriate method to diagnose GH deficiency, conclusions on the effects on the GH axis cannot be drawn (75). Finally, in a report of 22 patients with chronic pain on oral opioids, serum IGF-I levels and glucagon-stimulated peak GH were not different from a control group of 15 subjects on non-opioid analgesia (137).

The GH axis has also been assessed in male heroin addicts (138). ITT was performed in four groups: 12 controls (all denied opioid use), 19 methadone-treated patients, 16 heroin addicted patients, and 12 former heroin addicts. The mean of peak GH was similar in all groups. However, 26% of the methadone treated group and 31% of the heroin addicted people exhibited abnormal GH peak on the ITT (138).

Another remarkable aspect of the possible interplay between opioids and GH that deserves attention involves ghrelin which stimulates GH release via the growth hormone secretagogue receptor type 1a (GHS-R1a). Preclinical studies suggest that ghrelin/GHS-R1a has a significant role in the reinforcing effects of opioids, and in preclinical addiction studies, GHS-R1a antagonism decreased the opioid

reward and the symptoms of craving for opioids (139). Ghrelin/GHS-R1a involvement in opioid addiction in humans and the possible benefits of a ghrelin antagonist remain to be elucidated.

Clinical significance

Taken together, the scarce published literature suggests that long-term use of opioids may compromise the GH axis. Opioid type, dose and route of administration are potential parameters influencing these effects and more methodologically robust studies would provide definitive conclusions.

Interestingly, in a case report of a 61-year-old man with chronic intercostal neuropathic pain managed with methadone, GH treatment with daily doses up to 0.3 mg led to improvement of cognitive capacity (140). The patient had developed impaired cognitive function as a possible complication of opioid long-term treatment and had been diagnosed with GH deficiency by a GHRH-arginine test.

Overall, currently, evidence underpinning recommendations for monitoring, diagnosis, and management of GH deficiency in patients on long-term opioids users are not possible. However, health-care professionals prescribing opioids need to be aware of their possible negative effects on the GH axis.

Prolactin

Effects of opioids

Opioids can cause hyperprolactinemia most likely by inhibiting the hypothalamic dopamine production (49,141).

Early studies have shown that acute administration of opioids increases prolactin (PRL) secretion, an effect that is reversed by dopamine agonists (142). These actions have been confirmed in subsequent studies with various types of opioids (143-146).

In a systematic review of six studies evaluating the long-term effects of opioids on PRL secretion amongst 346 patients with either cancer pain or non-cancer pain, the results were variable (72). In a report of 14 patients on opioids for cancer pain, serum PRL levels were above the reference range in six (43%) of them. PRL values seemed to increase with higher opioid doses, but this finding did not reach statistical significance, possibly due to the small sample size (75). In a group of 39 non-cancer pain patients on oral opioids, hyperprolactinemia was detected in approximately 40% of them (118). In addition, high PRL levels were found in two out of 20 (10%) non-cancer pain patients on intrathecal morphine (147). In contrast, hyperprolactinemia has not been confirmed in other studies involving non-cancer pain patients on intrathecal or oral opioids (74,136,148), although in one of them, PRL was higher (but still within normal range) in men in the opioid treated group, as compared with the controls (148). In a report of 40 patients on oral or transdermal opioids for chronic pain (for more than 6 months on MME dose ≥ 25 mg), no difference in the prolactin values was found when compared with 25 controls not taking opioids (116). Finally, similar PRL levels were found between 21 non-cancer pain patients on oral opioids and a control group of 15 non-cancer pain patients on non-opioid analgesia; hyperprolactinemia was detected only in one case of the control group (137).

Higher PRL concentrations (as compared with healthy people) and hyperprolactinemia have been reported in people addicted to heroin and opium smokers (149). In patients on methadone or buprenorphine maintenance treatment, PRL has been found reduced, similar or higher when compared with controls (91,150,151).

Overall, based on the published literature, acute administration of opioids increases serum PRL, whereas in patients on long-term opioids, PRL has been found normal or elevated.

Clinical significance

Diagnostic approach and management

Given that hyperprolactinemia can cause hypogonadism and galactorrhea, health-care professionals prescribing opioids need to be aware of this possibility and enquire about these manifestations (152). Serum PRL should be measured only in the context of a suggestive clinical picture, and diagnostic steps should be in accord with published guidelines taking also into account the impact of potential confounding factors (e.g., pain, stress) (152).

It has been proposed that if hyperprolactinemia is confirmed, discontinuation or reduction of the opioid dose (or possibly use of an alternative opioid) need to be considered and PRL levels be re-evaluated (70). If these are not feasible, no other secondary etiology of hyperprolactinemia is identified, or the onset of the signs and symptoms do not coincide with the initiation of the opioid therapy, a magnetic resonance imaging of the sella turcica will be necessary to rule out a prolactinoma or other lesions compressing the pituitary stalk (49).

Therapeutic interventions are not needed in asymptomatic patients or if galactorrhea is not bothersome (49). Symptomatic patients who cannot discontinue or amend their opioid therapy could be treated with

a dopamine agonist, usually cabergoline. Alternatively, gonadal hormone replacement could be prescribed for hypogonadism (49).

DRAFT - DO NOT QUOTE OR CITE

Hypothalamo-pituitary-thyroid axis

Effects of opioids

Acute administration of opioids in healthy volunteers has led to increased serum TSH concentrations (153,154).

Studies with dynamic thyrotropin-releasing hormone (TRH) test performed in chronic opioid users have demonstrated conflicting results with enhanced (118), blunted (136,155,156) or no (74,157) thyroid-stimulating hormone (TSH) response. Furthermore, basal TSH values have been reported to be lower (149,153), unaffected (74,116,136,155-160) or increased (118,161). Variable results have also been described for the peripheral thyroid hormones; thyroxine (T4) levels were unaffected (74,118,136,159,160) or decreased (116,160-162) and triiodothyronine (T3) levels were increased (74,159) or unaffected (160) or decreased (161). T3 resin uptake, reflecting thyroid hormone binding globulin (TBG), has been reported to be lower (159).

A data mining study from the US Food and Drug Administration Adverse Event Reporting System (FAERS) demonstrated that in patients with OUD, there was positive association with hyperthyroidism (adjusted odds ratio 1.46, 95%CI: 1.43-1.49) and hypothyroidism (adjusted odds ratio 1.45, 95%CI: 1.42-1.48) (162). Whether this finding is truly significant, the potential pathophysiological mechanisms and possible links between opioids and auto-immune thyroid disease require further clarification.

Clinical significance

The data on the impact of opioids on thyroid function in humans are inconsistent. Overall, though, it can be speculated that long-term opioid use does not cause clinically relevant thyroid dysfunction, and, at present, there is no evidence supporting screening of thyroid disorders in these patients.

Arginine vasopressin

Effects of opioids

Earlier studies had suggested that morphine induces an antidiuresis in several species, including man, by stimulating the release of arginine vasopressin (AVP) (163,164). It has also been shown that intravenous or epidural morphine administered to children undergoing surgery had led to an increase in AVP levels (165). Fentanyl increased AVP release in both surgical patients (166) and healthy volunteers (167). Accordingly, opioids are mentioned in review articles as inducing the syndrome of inappropriate antidiuresis (SIAD) (168), an effect initially considered to be due to a direct enhancement of AVP release (169). The only opioid, however, that has been associated with hyponatremia in case reports is tramadol, a weak opioid analgesic with the hyponatremia usually occurring within the first 30 days after initiation of this drug (170-173). In a UK population-based cohort study, the use of tramadol was related with a two-fold increased risk of hospitalization for hyponatremia when compared to codeine; the authors postulated that hyponatremia developed due to stimulation of AVP and/or through inhibition of central re-uptake of serotonin, leading to increased AVP release and SIAD (174). According to the Beers criteria for Potentially Inappropriate Medication Use in Older Adults, tramadol should, therefore, be used with caution in older adults and sodium levels need to be monitored when starting this drug or when increasing its dose (175,176). It should be noted, however, that the results of many of these studies are difficult to be interpreted, as changes in patients' hemodynamic status or side effects of opioids (such as hypotension or nausea) which could secondarily stimulate AVP release, were not controlled.

In contrast to the above hypothesis, others have proposed that the primary effect of opioids is to suppress rather than stimulate AVP release. DAMME, a long-acting analogue of met-enkephalin, produced water diuresis in dehydrated or in hypertonic-saline-loaded subjects by inhibiting AVP release (177); in this study, the effect of opioids on AVP seemed to be dependent on the hydration status, as DAMME infused into water-deprived healthy volunteers produced diuresis which was attenuated by naloxone, whereas when subjects were hydrated, there was no change in water excretion (177). The inhibiting effect of

DAMME on AVP has been confirmed in further reports (178,179). It has also been shown that acute administration of the kappa-opioid receptor agonist asimadoline in normal subjects at doses 1, 5 and 10 mg reduced plasma AVP during 2.5% saline infusion in 6 out of 8 participants receiving the highest dose of 10 mg, as compared to placebo (180). Finally, opiate-dependent patients on diacetylmorphine (heroin) maintenance treatment had reduced AVP levels compared with controls (181). Thus, it may be speculated that the antidiuretic effect of opioids observed in some studies, as well as the occurrence of hyponatremia may not be mediated via AVP. Indeed, in a case report of a 19-year-old female with complete AVP deficiency since the age of 6 years, the patient developed hyponatremia upon treatment with hydrocodone offered after the extraction of her wisdom teeth (182). In this case, an increase of AVP levels could not explain the hyponatremia, as she was not able to produce AVP due to her AVP deficiency.

It is also worth looking at data on copeptin, a 39 amino acid long glycosylated peptide with a leucine-rich core region, which derives from the precursor protein Pre-Pro-Vasopressin together with AVP and Neurophysin II (183). In recent years, copeptin has emerged as a surrogate marker for AVP since, in contrast to AVP, it can be measured reliably in clinical routine by commercially available assays with high-standard technical performance. A strong correlation between plasma AVP (when measured with a well-established radio-immunoassay) and copeptin levels has been demonstrated in healthy volunteers (184), and, therefore, copeptin measurement could be an alternative marker to elucidate the effects of opioids on AVP. However, only very limited data on the effect of exogenous opioids on copeptin are available to date. A recent study measured copeptin levels before, during and after electrophysiological stimulation in healthy volunteers, initially prior to and then during opioid delivery; after administration of opioids, there was no visible effect on copeptin values (185).

Clinical significance

Based on the published literature, the effects of exogenous opioids on AVP secretion in humans are inconclusive and their clinical significance is unclear. The conflicting results might be due to

discrepancies in fluid/hydration status, as well as due to side effects of opioids leading to non-osmotically stimulated AVP. Given the finding that some opioids such as DAMME inhibit AVP release and induce diuresis in dehydrated subjects, careful hydration of patients receiving opioids may need to be considered. Furthermore, sodium monitoring is of importance in patients on tramadol, especially during the first 30 days of this treatment and this agent should be used with caution in older adults.

DRAFT - DO NOT QUOTE OR CITE

Oxytocin

Effects of opioids

Oxytocin receptors are present in the uterus, consistent with its known role in parturition, and in diverse other tissues. Oxytocin has a key role in the regulation of socio-emotional functioning, including attachment and pair bonding, fear extinction, emotion recognition, and empathy (186-189). In line with this, it may act as a stress-buffering hormone and influence the hypothalamic-pituitary-adrenal axis, as there was a trend toward attenuated cortisol levels by oxytocin (190). Besides the release from axonal terminals, there is dendritic release of oxytocin into the central extracellular space with local action and direct projection to other brain regions, including the amygdala, hippocampus, and brainstem, where it acts as a neurotransmitter (191). Specifically, over the past decades, it has been hypothesized that the prosocial effect of oxytocin might be secondary to an anxiolytic effect that involves modulation of amygdala responsivity, e.g., to threatening stimuli of different social valence (faces and scenes) (192).

Studies on the effects of opioids on oxytocin in humans have primarily focused on women, particularly investigating the effect of analgesics during pregnancy and labor. It has been shown that morphine and naloxone have no impact on oxytocin levels during pregnancy (193) and that morphine administration to the mother has no effect on fetal oxytocin levels (194). However, during the first stage of labor, morphine inhibited plasma oxytocin levels, whereas naloxone had no effect (195). Furthermore, after delivery, morphine inhibited the rise in oxytocin secretion in breast-feeding women but the amount of breast milk was not measured in this study (196). Similarly, other opioid analgesics, such as fentanyl, did not suppress oxytocin levels during pregnancy, but inhibited them during the onset of labor (197,198). Systemic opioids have a long tradition in analgesia during labor; in line with the inhibiting effect of opioids on oxytocin, in a non-randomised study, women offered opioid analgesia had longer labor and more need for oxytocin augmentation compared to controls not on labor analgesia (199). Women treated with intravenous opioids did not show oxytocin augmentation, as compared to those on epidural analgesia (200).

An intriguing aspect in the interplay between opioids and oxytocin is the effect of this hormone on opioid addiction and withdrawal. A randomized, double-blind clinical trial in heroin-dependent subjects showed that acute oxytocin treatment reduced craving and withdrawal scores, without, however, a significant effect on anxiety (201). It has been proposed that social attachment (the emotional bond or connection that individuals form with others, typically within social groups such as families, friends, or communities) seems to protect against addiction (202). It has, therefore, been postulated that oxytocin, through interaction with the central dopaminergic, serotonergic, and endogenous opioid system, could increase resilience against addiction by facilitating the process of social bonding (203).

Clinical significance

Data on the impact of opioids on oxytocin in humans outside pregnancy or labor are lacking. The clinical relevance of the finding that administration of opioids during labor might lead to higher need for oxytocin augmentation (to facilitate labor and birth) requires to be further investigated and also be correlated with labor outcomes (e.g., offering caesarean section, time to delivery, and risk of adverse perinatal outcomes). The proposed benefits of oxytocin treatment on opioid addiction and withdrawal are promising but they necessitate further validation before they are applied in clinical practice.

Bones

Effects of opioids

Exogenous opioids exert detrimental effects on bone homeostasis and metabolism and are associated with low bone mineral density (BMD) and an increased fracture risk. There are three main hypotheses that could explain the augmented fracture risk in chronic opioid users: bone loss caused by direct opioid effects on bone remodeling; increased risk of falls because of CNS effects; and chronic opioid-induced hypogonadism and its impact on bone metabolism (204).

Opioids and bone remodeling/bone healing

Data deriving from *in vitro* experiments on animal and human cells, as well as in animal models indicate that opioids can reduce BMD by directly influencing osteoblast activity (205-210). Pivotal experiments from two decades ago confirmed the expression of three types of opioid receptors, namely MOR, DOR, and KOR in the human osteoblast-like cell line MG-63 via RT-PCR and immunohistochemistry (208). Treatment with high concentrations of the MOR agonist morphine significantly inhibited the expression of osteocalcin, a marker of osteoblast activity, without influencing the secretion of alkaline phosphatase (208). In addition, treatment with the opioid antagonist naloxone reversed the effect of morphine with regards to osteocalcin expression (208). These results were replicated in a murine model, where treatment with morphine inhibited mineral nodule formation (207), and in human bone marrow-derived mesenchymal stem cells (MSCs), where morphine altered the ability of MSCs to differentiate into adipocytes or osteoblasts (206). Conversely, naltrexone, an antagonist of canonical opioid receptors and opioid growth factor receptor (OGFR), administered daily intraperitoneally to mice for 28 days increased their femoral bone mass, bone formation ratio, and osteoblast number/bone surface values when compared to control animals (210). Two additional experimental studies indicated a pro-osteogenic effect of naltrexone in a murine model of osteoporosis induced by liver cirrhosis (211,212). Thus, the findings of these preclinical studies suggest a pro-osteogenic role of opioid receptors

antagonists, although this will need to be validated in humans. Moreover, maternal consumption of morphine in rats led to impairment of longitudinal bone growth, as well inhibition of protein expression of chondrogenic and osteogenic markers in their offspring (209). A recent murine study using continuous morphine delivery via subcutaneous osmotic minipumps for 25 days indicated a sex difference in the bone outcome, where male mice treated with morphine had reduced trabecular bone volume and trabecular BMD in the distal femur compared with vehicle, while bone microarchitecture had not changed in females after morphine treatment (205). Of interest, this study also identified potential mediating miRNAs in the suppression of bone formation in male mice (205). Pregnant women addicted to heroin and cocaine displayed low serum levels of osteocalcin, a finding also collaborated in the umbilical arteries of their newborns, which also suggests toxic effects of these drugs on osteoblasts (213). Although potent opioid agonists exert negative effects on osteogenesis, the impact of opioids with lower affinity for the classic opioid receptors on bone appears to be less potent. For example, chronic use of tramadol, which has a lower affinity for MOR, led to a less pronounced osteoporotic phenotype in adult female rats compared to morphine or fentanyl (214). Furthermore, some interesting data have emerged on remifentanyl, an ultra-short acting opioid mainly used in the setting of peri-operative anaesthesia or analgesia in the intensive care unit (215). Under hypoxia-reoxygenation conditions, remifentanyl preconditioning enhanced the cell viability and maturation of human fetal osteoblasts (216), while in the pre-osteoblast C2C12 murine cell line, it increased osteoblast differentiation by upregulation of the expression of two key osteogenic transcription factors Runt-related transcription factor 2 (Runx2) and osterix (217).

The effects of opioids on osteoclasts are less well described. A number of studies have reported indirect effects on osteoclastogenesis and osteoclast function through modulation of osteoblast-derived factors. For example, tramadol was shown to enhance osteoprotegerin (OPG) synthesis in osteoblasts, thus indirectly suppressing the formation and function of osteoclasts (218,219). On the other hand, some evidence of direct effects of opioids on osteoclasts exists. In this vein, remifentanyl promoted the differentiation of murine pre-osteoclasts and induced their maturation in one experimental setting (220), although the exact opposite effects, namely reduction of the number and size of osteoclasts and the

formation of Tartrate-resistant acid phosphatase (TRAP)-positive multinuclear osteoclasts by remifentanyl, have also been described (221). In humans, higher serum concentrations of the bone resorption marker β C-terminal telopeptide (CTX) were reported in male heroin users compared to healthy volunteers (222), a finding replicated in men receiving opioid substitution therapy (223). Conversely, no up-regulation of CTX was found in another cohort of men on methadone substitution therapy (224).

Traditional opioids have been found to impair bone healing in some animal models of bone trauma (225). In a rat fracture model, morphine administration inhibited callus strength at 8 weeks (226), while fentanyl caused immature spinal fusion in a rabbit model (227). In contrast, low doses of naloxone enhanced bone mineralization and callus formation of a cortical defect induced in a sheep model (228).

In conclusion, data on the impact of opioids on bone remodeling and bone healing are discordant (225). Potential limitations for preclinical experiments appear to be different doses and duration of opioid application. Nonetheless, it appears that different opioid categories, also depending on their affinity for the respective opioid receptors, exert differential effects on bone cells. Moreover, most studies suggest differential effects on humans and preclinical models.

Clinical significance

Opioid prescription in the elderly population and bone health

Pain is common in elderly patients, being reported in up to 50% of the elderly population and up to 80% of care facilities inhabitants (229). Chronic pain is a limiting factor in daily activities and is linked to deconditioning, frailty, increased risk of accidents and polypharmacy (229). As the population ages, next to the burden to the individual, pain is bound to have considerable economical and societal impact (230). Thus, the associations between current comorbidities, clinical presentation and analgetic prescribing in elderly patients is a subject of interest.

In a large cohort of elderly patients admitted in a tertiary care hospital in Australia, analgesics were prescribed in 69% of them (more commonly in women), while opioids were prescribed in 34% of analgesic users (231). Female sex was associated with an increased likelihood of opioid prescription, with oxycodone being the most prescribed opioid (56%) (231). Another study investigating the prevalence and indications for daily opioid use among home care clients reported prevalence of 9.3% in opioid use for long periods of time (median duration of use was 357 days before study entry, and opioids were mostly continued during the follow-up year) (232). Most common indications for prescription were musculoskeletal disorders, i.e., vertebral osteoporotic fractures, degenerative spinal disorders, and osteoarthritis, with only 3.2% of the study population using an opioid for cancer-related pain (232). Similarly, a population-based cohort study of opioid-naïve community-dwelling US citizens reported continuous opioid use in 6.4% of patients one year after sustaining an index hip fracture (233). Of interest, a study assessing opioid use in the immediate 14 day-period post-hip fracture in older nursing home residents found that opioid use in that period was associated with a lower likelihood of death and functional decline (234). Regarding vertebral fractures, which are often associated with severe pain, vertebral augmentation procedures, such as balloon kyphoplasty and vertebroplasty were associated with decreased or discontinued opioid prescription fills in a significant proportion of 8.845 patients undergoing these procedures (235). In this vein, a second smaller study in a tertiary academic center confirmed decrease in pain scores, daily morphine consumption and improvement in functional status after kyphoplasty (236).

Bone mass in patients on opioids

Long-term opioid therapy is a risk factor for low BMD, as has been shown in various groups of patients. A prospective longitudinal nation-wide study of mid-life women assessed the impact of different analgesics [acetaminophen, nonsteroidal anti-inflammatory drugs (NSAIDs) and opioids] on bone mass and reported a more pronounced BMD decline over 5 years in opioid users (237). In contrast, an older

Danish cohort study with a 10-year follow-up period had failed to identify such a trend, albeit the number of opioid-users in that study was very small (238). In a small observational study comprising 20 males receiving long-term intrathecal opioid treatment for non-malignant bone pain, 17 patients displayed biochemical hypogonadism, while 10 of hypogonadal patients had osteopenia or osteoporosis (147). Of interest, in a study evaluating the effect of testosterone substitution in a small cohort of male patients developing hypogonadism under intrathecal opioid treatment, restoration of gonadal status and BMD after two years of testosterone therapy was reported (239). Another study including 12 men and 14 premenopausal women treated with oral opioids for chronic pain for at least one year found high prevalence of hypogonadism in men (75%), while only 21% of the women reported oligo- or amenorrhea indicating hypogonadism; osteopenia was detected in 50% of men and 21% of women, suggesting a potential sexual dimorphism in long-term effects of opioids regarding gonadal status and bone health (158). Lastly, in a cross-sectional analysis of Third National Health and Nutrition Examination Survey (NHANES III), opioids were associated with significantly reduced BMD after adjustment for several common causes of secondary osteoporosis (240).

A number of studies have investigated bone health of patients on opioid substitution therapy due to either previous addiction or chronic pain. A Swiss cross-sectional study including 144 men receiving opioid substitution therapy with either methadone (69%) or morphine (25%) or buprenorphine (6%) reported a decreased BMD compared to age- and BMI-matched healthy controls, predominantly on the axial skeleton in 73% of participants (223). In all age groups BMD was significantly lower than in age- and BMI- matched controls, while free testosterone was shown to be an independent predictor of bone mass (223). Of note, another cross-sectional study including 83 patients (48 men, 35 women) on methadone maintenance therapy (MMT) for a median of 11 years confirmed lower testosterone levels in men on MMT compared to controls and reported mean BMD values at each skeletal site approximately 10% below the population mean in men, albeit not in women (224). Furthermore, a small case-control study including young women on MMT compared with age- and BMI-matched controls found only marginally lower BMD at the hip of these subjects with no differences in other skeletal sites, despite a largely unhealthy lifestyle in patients (high alcohol and

nicotine intake) (241). A previous cross-sectional study in a mixed population (36% men) had described decreased BMD (compared to a gender and race/ethnicity-matched, young adult reference population) in 83% of people enrolled in a MMT program and identified male gender, low BMI, heavy alcohol use and HIV infection as significant predictors of low BMD; however, sex steroids were not evaluated in this study (242). In this vein, a prospective study aiming to investigate the combined effects of HIV infection and MMT in a sample of middle-aged women reported similar rates of decline in BMD between HIV-infected and HIV-uninfected women, but also indicated that MMT was associated with bone loss (243). A recent systematic review assessing all the above studies on patients on opioid substitution therapy concluded that this population should be screened for low bone mass and that risk factors significantly associated with low BMD were male gender, low BMI, low testosterone levels, methadone or heroin use and longer duration of heavy alcohol use (244) .

In conclusion, low BMD is a long-term side-effect of opioids, and especially in male patients, it appears to be pathophysiologically connected to hypogonadism.

Fracture risk in patients on opioids

Use of exogenous opioids has been associated with an increased risk of fractures in several meta-analyses (245-248). Using mostly cohort and case-controls studies, these meta-analyses have reported an increased risk for hip fractures (245), falls (247) and major osteoporotic fractures (246,248) in opioid-treated patients.

As previously stated, the increased fracture risk among individuals with chronic opioid use is attributed to three main reasons: reduced BMD caused by direct opioid effects on bone metabolism, chronic opioid-induced hypogonadism, and increased risk of falls caused by CNS side-effects, including sedation, gait disorders and lack of concentration. The first two mechanisms have been elaborated above. Regarding CNS side-effects, it has been shown that the rates of hospital visits for falls and fractures is disproportionally high at the initiation phase of opioid treatment, a trend not seen with other

analgesics, such as NSAIDs (249). Of note, such an increase in falls and fractures is accentuated in new opioid users with cognitive impairment, as seen in a study of patients with Alzheimer's disease (250). Conversely, some data including time-to-event analyses, highlight a high incidence of falls and fractures in long-time opioid users, as reported for example in non-tramadol opioid users with osteoarthritis experiencing fractures in a median time of 14 months after opioid initiation (251). Thus, it appears that patients with continuous opioid use remain at high risk for falls and fracture.

Diagnostic approach and management

Acknowledging the fact that majority of data on skeletal effects of opioids derives from observational studies characterized by considerable heterogeneity, various confounders (e.g., poor global nutrition, acidosis, low calcium and vitamin D intake, low body weight, weight loss, smoking, alcohol, use of additional drugs seizures, reduced muscle mass, inactivity, chronic medical condition/malignancy) and different follow-up periods, it appears indisputable that opioid use is associated with detrimental skeletal consequences. Thus, measurement of BMD by Dual-energy X-ray absorptiometry (DXA) and evaluation of additional osteoporosis risk factors is indicated in long-time opioid users, especially in the presence of hypogonadism. Careful assessment of the necessity of chronic opioid treatment and de-escalation/discontinuation of their use is of major importance. In case of osteopenia/osteoporosis, management should be tailored to current osteoporosis guidelines. Scarce data exist on the effects of sex hormone substitution on bone health in hypogonadal patients on opioids. A small observational study of male hypogonadal patients on intrathecal opioids for non-cancer pain who received testosterone substitution corrected the decreased testosterone levels and improved BMD (239). Regular exercise has effects on bone density, size, and shape, resulting in substantial improvements in mechanical strength (252), but data on the effect of exercise interventions on the bone quality of opioid-dependent individuals are limited. An 8-month aerobic gymnastics training program in young opioid-addicted women failed to show any effects on bone quality, as assessed by quantitative ultrasound (253). Additional large well-designed studies investigating such approaches in individuals on chronic opioid use will underpin further guidance.

Summary of key points on actions of exogenous opioids on endocrine system in humans

- Opioids inhibit the hypothalamo-pituitary-gonadal axis and can cause hypogonadism.
- Opioids inhibit the hypothalamo-pituitary-adrenal axis; low basal morning cortisol levels and inadequate response of cortisol on dynamic testing have been reported; the significance of these effects and the occurrence of clinically relevant adrenal insufficiency is not fully elucidated.
- Acute administration of opioids has a stimulatory effect on GH secretion; data on the impact of long-term use of opioids on the GH axis are not consistent but cases with compromised response of GH on dynamic testing have been reported.
- Acute administration of opioids increases prolactin secretion. Chronic use of opioids can lead to hyperprolactinemia.
- Opioids have variable and inconsistent effects on the hypothalamo-pituitary-thyroid axis.
- Data on the effects of opioids on AVP secretion are inconclusive.
- Evidence of opioid actions on oxytocin derived from studies of women in labor suggests an inhibitory effect (opioids can inhibit plasma oxytocin levels during labor).
- Opioid use can have negative impact on bone homeostasis and metabolism; reduced BMD and increased risk of fracture have been reported; proposed implicated mechanisms for these negative sequelae include direct effects on bone remodeling, increased risk of falls and hypogonadism.

Gaps in knowledge

Despite the global opioid crisis, the gaps in knowledge around the actions of opioids on the endocrine systems, their clinical significance and their optimal management are wide. In Table 3, we summarise the most pertinent ones on the areas this Statement has covered which require careful consideration.

Table 3. Gaps in knowledge on the effects of exogenous opioids in humans on the endocrine systems covered in the Statement.

Hypothalamo-Pituitary-Gonadal axis
Data on type, route of administration and regimens of opioids associated with higher risk of hypogonadism in both males and females derived from studies adjusting for confounding factors. Impact of partial agonists (e.g., buprenorphine).
Impact of opioids on fertility in males and females.
Time frame of recovery of the hypothalamo-pituitary-gonadal axis after opioid discontinuation to inform management protocols.
Benefits and risks of gonadal hormone replacement in pre-menopausal females and in males with hypogonadism attributed to opioids from prospective adequately powered and of appropriate duration randomised trials; this also includes impact on pain sensitivity and on the effects on recovery of people on methadone maintenance therapy.
Hypothalamo-Pituitary-Adrenal Axis
Standardization and validation of diagnostic tests and criteria for opioid-induced adrenal insufficiency.
Prevalence of hypoadrenalism with different opioids, doses, and routes of administration.
Understand the individual susceptibility to the inhibitory effects of opioids (could differences in opioid metabolism combined with a possible opioid concentration threshold which inhibits the axis explain why some individuals are more susceptible to these effects?).
Identify patients at risk of developing adrenal insufficiency whilst on opioids.
Clinical significance and optimal management approach of opioid-induced adrenal insufficiency (need for daily physiological glucocorticoid replacement or cover with glucocorticoids in case of stress?).
Time course of the reversal of the changes in the axis following withdrawal or reduction of opioid dose to inform management protocols.
GH axis
Clear understanding of the effects of chronic use of opioids on the GH axis (with different opioids, doses, and routes of administration).
Clinical significance of compromised GH axis in patients on opioids for cancer or non-cancer pain and in people with opioid use disorder and potential benefits/risks of treatment with GH replacement.
PRL
Types, doses, and route of administration of opioids etiologically related with clinically significant hyperprolactinemia.
Benefits and risks of dopamine agonist treatment in patients with opioid-induced hyperprolactinemia.
Consequences of untreated long-term opioid-induced hyperprolactinemia.
Hypothalamo-pituitary-thyroid axis
True effects of opioids on thyroid function and their clinical relevance.
Implications of opioids in the development of autoimmune thyroid disease.
Arginine vasopressin

True impact of different opioids on AVP release after adjusting for confounding factors and considering hydration status of the patients.
Effects of opioids on copeptin as a surrogate marker of AVP secretion.
Mechanisms leading to hyponatremia in patients treated with opioids (AVP-dependent or AVP-independent?).
Oxytocin
Effects of opioids on oxytocin in males and in non-pregnant women.
Clinically relevant impact of opioids offered as analgesics on labor outcomes.
Role of oxytocin in the treatment of opioid addiction and withdrawal.
Bone health
Differential effects of various opioid agonists on bone homeostasis and factors related with them.
Role of opioid antagonists in increasing BMD and possible treatment for osteoporosis.
Sexual dimorphism of opioid effects in bone homeostasis and its significance.
Impact of gonadal hormone replacement on improving/reversing reduced BMD and on risk of fracture in long-term opioid users.

GH growth hormone; PRL prolactin; AVP arginine vasopressin; BMD bone mineral density.

Conclusions/Future directions

This Scientific Statement presents an overview of the landscape on research relating to the effects of exogenous opioids on various components of the human endocrine system. It specifically describes clinically relevant data on the actions of opioids on the hypothalamo-pituitary system and on bone health. It also identifies areas requiring further research that will enhance our understanding on the consequences of the available opioids and will provide evidence guiding the optimal diagnostic approach and management of their endocrine adverse sequelae. These are additionally mandated by the opioid pandemic we are witnessing and will open perspectives for exciting and impactful future research.

The true prevalence and the risk of opioid-induced hypogonadism in both males and females in various patient cohorts needs to be clarified in methodologically sound studies with adjustment for confounding factors (e.g., pain, chronic stress, co-morbidities). Of particular importance is the identification of the types and doses of opioids associated with high potential of causing hypogonadism and the time frame for restoration of the hypothalamo-pituitary-gonadal axis after their cessation or dose reduction. Randomised controlled adequately powered and long duration trials are required to assess the effects of gonadal hormone replacement not only on known manifestations/consequences of hypogonadism but also on aspects related to pain management and recovery of people on methadone maintenance therapy. The paucity of data on females highlights the need for studies in women which will also look at fertility outcomes.

Research is needed to establish consensus diagnostic criteria for adrenal insufficiency in patients on opioids. Without them, future studies will not be directly comparable. Increasing use of liquid chromatography-tandem mass spectrometry for the measurement of cortisol should largely eliminate the cut-off variability observed using immunoassay technology, both for basal morning cortisol concentrations and for stimulation tests. More detailed pharmacokinetic, pharmacodynamic and pharmacogenomic studies need to be undertaken to improve our understanding of relationship between

opioid dose, plasma concentration and causation of secondary adrenal insufficiency (and also of other opioid-related endocrine effects), aiming to identify people who might potentially be at increased risk of developing opioid-induced adrenal insufficiency. The evidence behind recommendations to routinely treat patients all diagnosed with opioid-induced secondary adrenal insufficiency using standard physiological glucocorticoid replacement doses is weak and a well powered, double-blind randomised control trial comparing hydrocortisone *versus* placebo is required to definitively answer the question. In addition, the time course of hypothalamo-pituitary-adrenal axis recovery during the process of opioid withdrawal requires more detailed documentation in a longitudinal manner and whether administration of full replacement glucocorticoid doses during this process impedes or promotes recovery – symptomatically and functionally.

The scarce published literature suggests that long-term opioid use could possibly impact negatively on the GH axis in humans. Nonetheless, the existing evidence is not sufficient to elucidate this area and clinical recommendations are not possible. Methodologically robust studies, adequately powered and with appropriate controls are needed to reveal the true impact of various opioids and its clinical significance, as well as the benefits of testing the integrity of the GH axis and offering GH replacement in cases with biochemically confirmed GH deficiency and no contraindications for this treatment.

Long-term use of opioids has been associated with hyperprolactinemia, the true prevalence of which is not clear necessitating studies to clarify its frequency depending on the opioid, dose, and route of administration. The optimal management approach for the hyperprolactinemia including the use of dopamine agonists requires further research.

The effect of opioids on the hypothalamo-pituitary-thyroid axis require longitudinal studies looking at biochemical measures and clinical parameters reflecting thyroid dysfunction, acknowledging the baseline risk to encounter thyroid disease in general.

The effects of opioids on AVP in humans are inconclusive and their clinical significance is not well defined. Further studies providing insights on the interplay between opioids, AVP/copeptin secretion, and water balance, as well as the role of hydration status on the effects of these agents are needed. Other confounding factors will need to be considered in the design of these studies (e.g., influence of pain and nausea on non-osmotic AVP secretion).

The clinical relevance of the inhibitory action of opioids during labor necessitates further investigation. Additional studies are also needed to clarify the effects of opioids on oxytocin in males and in non-pregnant women and to explore and validate potential benefits of oxytocin in the treatment of opioid addiction and withdrawal.

Further research is needed to illuminate the different effects of various opioids on bone metabolism and whether these are characterised by sexual dimorphism. In addition, well-designed prospective studies accounting for confounding factors and of appropriate follow-up will elucidate the long-term benefits of hormonal replacement on BMD and facilitate the development of safe and cost-effective guidelines on the bone health of chronic opioid users.

Finally, studies investigating the awareness of clinicians (and particularly those prescribing opioids) and of patients on the endocrine consequences of these agents will identify gaps in the provision of care in these groups of patients and will offer additional insights on the implications of the opioid crisis on the health care system.

Disclaimer statement

The Endocrine Society develops Scientific Statements to explore the scientific basis of hormone-related conditions and disease, discuss how this knowledge can be applied in practice, and identify areas that require additional research. Topics are selected on the basis of their emerging scientific impact. Scientific Statements are developed by a Task Force of experts appointed by the Endocrine

Society, with internal review by the relevant Society committees and expert external reviewers prior to a comment period open to all members of the Society. Clinicians and other users should not consider this Scientific Statement inclusive of all proper approaches or methods or exclusive of others. It cannot guarantee any specific outcome, nor does it establish a standard of care. It is not intended to dictate the treatment of a particular patient. The independent judgment of health care providers and each patient's individual circumstances must determine treatment decisions. The Endocrine Society makes no warranty, express or implied, regarding this Scientific Statement and specifically excludes any warranties of merchantability and fitness for a particular use or purpose. The Endocrine Society shall not be liable for direct, indirect, special, incidental, or consequential damages related to the use of the information contained herein.

Funding: The authors did not receive support from any organization for the submitted work.

Disclosures: Niki Karavitaki: Speaker for Pfizer, Ipsen, HRA Pharma, Recordati Rare Diseases Consilient Health - Investigator for Pfizer, Ipsen, Shire - Scientific Advisory Board for Pfizer, Ipsen, Recordati Rare Diseases; Monica R. Gadelha: Speaker for Ipsen, Recordati Rare Diseases, Novo Nordisk - Investigator for Crinetics, Recordati Rare Diseases - Scientific Advisory Board for Crinetics, Recordati Rare Diseases, Novo Nordisk, Ipsen; Warrick Inder: Scientific Advisory Board for Recordati Rare Diseases; Elena Tsourdi: honoraria for lectures and advisory boards from Amgen, UCB, Alexion, Ascendis and educational grants from UCB; Maria Chiara Zatelli: Consultant Lilly - Grants from Savio, AAA, Sanofi, Sandoz - Investigator for Sandoz.

References

1. Brownstein MJ. A brief history of opiates, opioid peptides, and opioid receptors. *Proc Natl Acad Sci U S A*. 1993;90(12):5391-5393.
2. Opioid Overdose. <https://www.cdc.gov/drugoverdose/deaths/opioid-overdose.html>. Access February 24, 2024. Centers for Disease Control and Prevention.
3. O'Donnell J, Tanz LJ, Gladden RM, Davis NL, Bitting J. Trends in and Characteristics of Drug Overdose Deaths Involving Illicitly Manufactured Fentanyls - United States, 2019-2020. *MMWR Morb Mortal Wkly Rep*. 2021;70(50):1740-1746.
4. Jones CM, Bekheet F, Park JN, Alexander GC. The Evolving Overdose Epidemic: Synthetic Opioids and Rising Stimulant-Related Harms. *Epidemiol Rev*. 2020;42(1):154-166.
5. Weiss AJ, Elixhauser A, Barrett ML, Steiner CA, Bailey MK, O'Malley L. Opioid-Related Inpatient Stays and Emergency Department Visits by State, 2009–2014. Healthcare Cost and Utilization Project (HCUP) Statistical Briefs. Rockville (MD): Agency for Healthcare Research and Quality (US); 2006.
6. Provisional Drug Overdose Death Counts. <https://www.cdc.gov/nchs/nvss/vsrr/drug-overdose-data.htm>. Access February 24, 2024. Centers for Disease Control and Prevention. .
7. UNODC, World Drug Report 2022 (United Nations publication, 2022). Booklet 3 - Drug market trends of Cannabis and Opioids. United Nations Office on Drugs and Crime; 2022: https://www.unodc.org/unodc/en/data-and-analysis/wdr-2022_booklet-3.html.
8. Hedegaard H MA, Warner M. . Drug overdose deaths in the United States, 1999–2017. <https://www.cdc.gov/nchs/data/databriefs/db329-h.pdf>. *NCHS Data Brief*, no 329. 2018.
9. Warner M, Trinidad JP, Bastian BA, Minino AM, Hedegaard H. Drugs Most Frequently Involved in Drug Overdose Deaths: United States, 2010-2014. *Natl Vital Stat Rep*. 2016;65(10):1-15.
10. United States Dispensing Rate Maps. <https://www.cdc.gov/drugoverdose/rxrate-maps/index.html> . Access March 2, 2024. Vol March 2, 2024.
11. Cicero TJ, Ellis MS, Harney J. Shifting Patterns of Prescription Opioid and Heroin Abuse in the United States. *N Engl J Med*. 2015;373(18):1789-1790.
12. Cicero TJ, Ellis MS, Kasper ZA. Polysubstance Use: A Broader Understanding of Substance Use During the Opioid Crisis. *Am J Public Health*. 2020;110(2):244-250.
13. Schieber LZ, Guy GP, Jr., Seth P, Young R, Mattson CL, Mikosz CA, Schieber RA. Trends and Patterns of Geographic Variation in Opioid Prescribing Practices by State, United States, 2006-2017. *JAMA Netw Open*. 2019;2(3):e190665.
14. Chen Q, Larochelle MR, Weaver DT, Lietz AP, Mueller PP, Mercaldo S, Wakeman SE, Freedberg KA, Raphael TJ, Knudsen AB, Pandharipande PV, Chhatwal J. Prevention of Prescription Opioid Misuse and Projected Overdose Deaths in the United States. *JAMA Netw Open*. 2019;2(2):e187621.
15. Ciccarone D. The triple wave epidemic: Supply and demand drivers of the US opioid overdose crisis. *Int J Drug Policy*. 2019;71:183-188.
16. Ciccarone D. The rise of illicit fentanyls, stimulants and the fourth wave of the opioid overdose crisis. *Curr Opin Psychiatry*. 2021;34(4):344-350.
17. Strickland JC, Havens JR, Stoops WW. A nationally representative analysis of "twin epidemics": Rising rates of methamphetamine use among persons who use opioids. *Drug Alcohol Depend*. 2019;204:107592.
18. Argoff CE, Alford DP, Fudin J, Adler JA, Bair MJ, Dart RC, Gandolfi R, McCarberg BH, Stanos SP, Gudín JA, Polomano RC, Webster LR. Rational Urine Drug Monitoring in Patients Receiving Opioids for Chronic Pain: Consensus Recommendations. *Pain Med*. 2018;19(1):97-117.

19. Dowell D, Haegerich TM, Chou R. CDC Guideline for Prescribing Opioids for Chronic Pain - United States, 2016. *MMWR Recomm Rep*. 2016;65(1):1-49.
20. Chou R, Fanciullo GJ, Fine PG, Adler JA, Ballantyne JC, Davies P, Donovan MI, Fishbain DA, Foley KM, Fudin J, Gilson AM, Kelter A, Mauskop A, O'Connor PG, Passik SD, Pasternak GW, Portenoy RK, Rich BA, Roberts RG, Todd KH, Miaskowski C. Clinical guidelines for the use of chronic opioid therapy in chronic noncancer pain. *J Pain*. 2009;10(2):113-130.
21. National Quality Partners Playbook™: Opioid Stewardship National Quality Forum.
22. Pauly Jb PharmD B, Vartan Cm PharmD B, Brooks At PharmD B. Implementation and evaluation of an opioid overdose education and naloxone distribution (OEND) program at a Veterans Affairs Medical Center. *Subst Abus*. 2018;39(2):206-210.
23. RA P. Opioid Safety Initiative Requirements. Memorandum from VA Under Secretary for Health April 2, 2014. 2014.
24. Comprehensive Addiction and Recovery Act of 2016 Public Law 114-198, Signed July 22, 2016.
25. VA/DoD Clinical Practice Guideline for Opioid Therapy for Chronic Pain. February 2017.
26. Duensing K, Twillman R, Ziegler S, Cepeda MS, Kern D, Salas M, Wedin G. An Examination of State and Federal Opioid Analgesic and Continuing Education Policies: 2016-2018. *J Pain Res*. 2020;13:2431-2442.
27. World Drug Report 2020 (United Nations publication, Sales No. E.20.XI.6). Booklet 2 - Drug use and health consequences United Nations Office on Drugs and Crime; 2020.
28. Häuser W, Buchser E, Finn DP, Dom G, Fors E, Heiskanen T, Jarlbaek L, Knaggs RD, Kosek E, Krceviski-Škvarč N, Pakkonen K, Perrot S, Trouvin AP, Morlion B. Is Europe also facing an opioid crisis?-A survey of European Pain Federation chapters. *Eur J Pain*. 2021;25(8):1760-1769.
29. Report of the International Narcotics Control Board for 2022.
30. Diagnostic and statistical manual of mental disorders: DSM-5™ (5th ed.). . American Psychiatric Publishing, Inc.
31. Dela Cruz AM, Walker R, Pipes R, Wakhlu S, Trivedi MH. Creation of an algorithm for clinical decision support for treatment of opioid use disorder with buprenorphine in primary care. *Addict Sci Clin Pract*. 2021;16(1):12.
32. Marsden J, Tai B, Ali R, Hu L, Rush AJ, Volkow N. Measurement-based care using DSM-5 for opioid use disorder: can we make opioid medication treatment more effective? *Addiction*. 2019;114(8):1346-1353.
33. Substance Abuse and Mental Health Services Administration. (2023). Key substance use and mental health indicators in the United States: Results from the 2022 National Survey on Drug Use and Health (HHS Publication No. PEP23-07-01-006, NSDUH Series H-58). Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. <https://www.samhsa.gov/data/report/2022-nsduh-annual-national-report>.
34. Strang J, Volkow ND, Degenhardt L, Hickman M, Johnson K, Koob GF, Marshall BDL, Tyndall M, Walsh SL. Opioid use disorder. *Nat Rev Dis Primers*. 2020;6(1):3.
35. Florence C, Luo F, Rice K. The economic burden of opioid use disorder and fatal opioid overdose in the United States, 2017. *Drug Alcohol Depend*. 2021;218:108350.
36. James K, Jordan A. The Opioid Crisis in Black Communities. *J Law Med Ethics*. 2018;46(2):404-421.
37. Parlier-Ahmad AB, Pugh M, Jr., Martin CE. Treatment Outcomes Among Black Adults Receiving Medication for Opioid Use Disorder. *J Racial Ethn Health Disparities*. 2022;9(4):1557-1567.
38. Saloner B, Karthikeyan S. Changes in Substance Abuse Treatment Use Among Individuals With Opioid Use Disorders in the United States, 2004-2013. *Jama*. 2015;314(14):1515-1517.
39. Stahler GJ, Mennis J. Treatment outcome disparities for opioid users: Are there racial and ethnic differences in treatment completion across large US metropolitan areas? *Drug Alcohol Depend*. 2018;190:170-178.
40. Stein BD, Dick AW, Sorbero M, Gordon AJ, Burns RM, Leslie DL, Pacula RL. A population-based examination of trends and disparities in medication treatment for opioid use disorders among Medicaid enrollees. *Subst Abus*. 2018;39(4):419-425.

41. Wu LT, Zhu H, Swartz MS. Treatment utilization among persons with opioid use disorder in the United States. *Drug Alcohol Depend.* 2016;169:117-127.
42. Kariisa M, Davis NL, Kumar S, Seth P, Mattson CL, Chowdhury F, Jones CM. Vital Signs: Drug Overdose Deaths, by Selected Sociodemographic and Social Determinants of Health Characteristics - 25 States and the District of Columbia, 2019-2020. *MMWR Morb Mortal Wkly Rep.* 2022;71(29):940-947.
43. National Academies of Sciences, Engineering, and Medicine. 2019. Medications for opioid use disorder save lives. Washington, DC: The National Academies Press. Doi: <https://doi.org/10.17226/25310>.
44. Crotty K, Freedman KI, Kampman KM. Executive Summary of the Focused Update of the ASAM National Practice Guideline for the Treatment of Opioid Use Disorder. *J Addict Med.* 2020;14(2):99-112.
45. Vallejo R, Barkin RL, Wang VC. Pharmacology of opioids in the treatment of chronic pain syndromes. *Pain Physician.* 2011;14(4):E343-360.
46. Waldhoer M, Bartlett SE, Whistler JL. Opioid receptors. *Annu Rev Biochem.* 2004;73:953-990.
47. Maehle AH, Prüll CR, Halliwell RF. The emergence of the drug receptor theory. *Nat Rev Drug Discov.* 2002;1(8):637-641.
48. Raffa RB, Haidery M, Huang HM, Kalladeen K, Lockstein DE, Ono H, Shope MJ, Sowunmi OA, Tran JK, Pergolizzi JV, Jr. The clinical analgesic efficacy of buprenorphine. *J Clin Pharm Ther.* 2014;39(6):577-583.
49. Gadelha MR, Karavitaki N, Fudin J, Bettinger JJ, Raff H, Ben-Shlomo A. Opioids and pituitary function: expert opinion. *Pituitary.* 2022;25(1):52-63.
50. Grassin-Delyle S, Buenestado A, Naline E, Faisy C, Blouquit-Laye S, Couderc LJ, Le Guen M, Fischler M, Devillier P. Intranasal drug delivery: an efficient and non-invasive route for systemic administration: focus on opioids. *Pharmacol Ther.* 2012;134(3):366-379.
51. CDC's Guideline for Prescribing Opioids. Module 6: Dosing and Titration of Opioids: How Much, How Long, and How and When to Stop? <https://cdc.gov/drugoverdose/training/dosing/accessible/index.html>.
52. Schumacher MA BA, Naidu RK. Opioid Agonists & Antagonists. Basic & Clinical Pharmacology. 15th ed: McGraw Hill; 2021.
53. Al-Hasani R, Bruchas MR. Molecular mechanisms of opioid receptor-dependent signaling and behavior. *Anesthesiology.* 2011;115(6):1363-1381.
54. Dahan A, van der Schrier R, Smith T, Aarts L, van Velzen M, Niesters M. Averting Opioid-induced Respiratory Depression without Affecting Analgesia. *Anesthesiology.* 2018;128(5):1027-1037.
55. Bettinger J TN, Fudin J, Wegrzyn E, Schatman M. Fentanyl: Separating Fact from Fiction. *Pract Pain Manag.* 2018;18(5).
56. Mazák K, Noszál B, Hosztafi S. Advances in the Physicochemical Profiling of Opioid Compounds of Therapeutic Interest. *ChemistryOpen.* 2019;8(7):879-887.
57. Oldendorf WH. Lipid solubility and drug penetration of the blood brain barrier. *Proc Soc Exp Biol Med.* 1974;147(3):813-815.
58. Schaefer CP, Tome ME, Davis TP. The opioid epidemic: a central role for the blood brain barrier in opioid analgesia and abuse. *Fluids Barriers CNS.* 2017;14(1):32.
59. Volpe DA, McMahon Tobin GA, Mellon RD, Katki AG, Parker RJ, Colatsky T, Kropp TJ, Verbois SL. Uniform assessment and ranking of opioid μ receptor binding constants for selected opioid drugs. *Regul Toxicol Pharmacol.* 2011;59(3):385-390.
60. Webster L, Gudín J, Raffa RB, Kuchera J, Rauck R, Fudin J, Adler J, Mallick-Searle T. Understanding Buprenorphine for Use in Chronic Pain: Expert Opinion. *Pain Med.* 2020;21(4):714-723.
61. Böttcher B, Seeber B, Leyendecker G, Wildt L. Impact of the opioid system on the reproductive axis. *Fertil Steril.* 2017;108(2):207-213.
62. Petraglia F, Porro C, Facchinetti F, Cicoli C, Bertellini E, Volpe A, Barbieri GC, Genazzani AR. Opioid control of LH secretion in humans: menstrual cycle, menopause and aging reduce effect of naloxone but not of morphine. *Life Sci.* 1986;38(23):2103-2110.

- 1529 63. Vuong C, Van Uum SH, O'Dell LE, Lutfy K, Friedman TC. The effects of opioids and opioid
1530 analogs on animal and human endocrine systems. *Endocr Rev.* 2010;31(1):98-132.
- 1531 64. Wildt L, Sir-Petermann T, Leyendecker G, Waibel-Treber S, Rabenbauer B. Opiate antagonist
1532 treatment of ovarian failure. *Hum Reprod.* 1993;8 Suppl 2:168-174.
- 1533 65. Fountas A, Chai ST, Kourkouti C, Karavitaki N. MECHANISMS OF ENDOCRINOLOGY:
1534 Endocrinology of opioids. *Eur J Endocrinol.* 2018;179(4):R183-r196.
- 1535 66. Purohit V, Singh HH, Ahluwalia BS. Failure of methadone-treated human testes to respond to
1536 the stimulatory effect of human chorionic gonadotrophin on testosterone biosynthesis in vitro.
1537 *J Endocrinol.* 1978;78(2):299-300.
- 1538 67. Ragni G, De Lauretis L, Bestetti O, Sghedoni D, Gambaro V. Gonadal function in male heroin
1539 and methadone addicts. *Int J Androl.* 1988;11(2):93-100.
- 1540 68. Safarinejad MR, Asgari SA, Farshi A, Ghaedi G, Kolahi AA, Iravani S, Khoshdel AR. The
1541 effects of opiate consumption on serum reproductive hormone levels, sperm parameters,
1542 seminal plasma antioxidant capacity and sperm DNA integrity. *Reprod Toxicol.* 2013;36:18-
1543 23.
- 1544 69. Wersocki E, Bedson J, Chen Y, LeResche L, Dunn KM. Comprehensive systematic review of
1545 long-term opioids in women with chronic noncancer pain and associated reproductive
1546 dysfunction (hypothalamic-pituitary-gonadal axis disruption). *Pain.* 2017;158(1):8-16.
- 1547 70. Fountas A, Van Uum S, Karavitaki N. Opioid-induced endocrinopathies. *Lancet Diabetes*
1548 *Endocrinol.* 2020;8(1):68-80.
- 1549 71. Pelosi MA, Sama JC, Caterini H, Kaminetzky HA. Galactorrhea-amenorrhea syndrome
1550 associated with heroin addiction. *Am J Obstet Gynecol.* 1974;118(7):966-970.
- 1551 72. de Vries F, Bruin M, Lobatto DJ, Dekkers OM, Schoones JW, van Furth WR, Pereira AM,
1552 Karavitaki N, Biermasz NR, Zamanipoor Najafabadi AH. Opioids and Their Endocrine Effects:
1553 A Systematic Review and Meta-analysis. *J Clin Endocrinol Metab.* 2020;105(3):1020-1029.
- 1554 73. Daniell HW. Opioid endocrinopathy in women consuming prescribed sustained-action opioids
1555 for control of nonmalignant pain. *J Pain.* 2008;9(1):28-36.
- 1556 74. Abs R, Verhelst J, Maeyaert J, Van Buyten JP, Opsomer F, Adriaensen H, Verlooy J, Van
1557 Havenbergh T, Smet M, Van Acker K. Endocrine consequences of long-term intrathecal
1558 administration of opioids. *J Clin Endocrinol Metab.* 2000;85(6):2215-2222.
- 1559 75. Merdin A, Merdin FA, Gündüz Ş, Bozcuk H, Coşkun H. Opioid endocrinopathy: A clinical
1560 problem in patients with cancer pain. *Exp Ther Med.* 2016;11(5):1819-1822.
- 1561 76. Strasser F, Palmer JL, Schover LR, Yusuf SW, Pisters K, Vassilopoulou-Sellin R, DeGracia B,
1562 Willey JS, Bruera E. The impact of hypogonadism and autonomic dysfunction on fatigue,
1563 emotional function, and sexual desire in male patients with advanced cancer: a pilot study.
1564 *Cancer.* 2006;107(12):2949-2957.
- 1565 77. McWilliams K, Simmons C, Laird BJ, Fallon MT. A systematic review of opioid effects on the
1566 hypogonadal axis of cancer patients. *Support Care Cancer.* 2014;22(6):1699-1704.
- 1567 78. Rajagopal A, Vassilopoulou-Sellin R, Palmer JL, Kaur G, Bruera E. Symptomatic
1568 hypogonadism in male survivors of cancer with chronic exposure to opioids. *Cancer.*
1569 2004;100(4):851-858.
- 1570 79. Aloisi AM, Aurilio C, Bachiocco V, Biasi G, Fiorenzani P, Pace MC, Paci V, Pari G, Passavanti
1571 G, Ravaioli L, Sindaco G, Vellucci R, Ceccarelli I. Endocrine consequences of opioid therapy.
1572 *Psychoneuroendocrinology.* 2009;34 Suppl 1:S162-168.
- 1573 80. Mendelson JH, Mello NK. Plasma testosterone levels during chronic heroin use and protracted
1574 abstinence. A study of Hong Kong addicts. *Clin Pharmacol Ther.* 1975;17(5):529-533.
- 1575 81. Finch PM, Roberts LJ, Price L, Hadlow NC, Pullan PT. Hypogonadism in patients treated with
1576 intrathecal morphine. *Clin J Pain.* 2000;16(3):251-254.
- 1577 82. Mendelson JH, Mendelson JE, Patch VD. Plasma testosterone levels in heroin addiction and
1578 during methadone maintenance. *J Pharmacol Exp Ther.* 1975;192(1):211-217.
- 1579 83. Mendelson JH, Inturrisi CE, Renault P, Senay EC. Effects of acetylmethadol on plasma
1580 testosterone. *Clin Pharmacol Ther.* 1976;19(3):371-374.
- 1581 84. Policola C, Stokes V, Karavitaki N, Grossman A. Adrenal insufficiency in acute oral opiate
1582 therapy. *Endocrinol Diabetes Metab Case Rep.* 2014;2014:130071.

85. Woody G, McLellan AT, O'Brien C, Persky H, Stevens G, Arndt I, Carroff S. Hormone secretion in methadone-dependent and abstinent patients. *NIDA Res Monogr*. 1988;81:216-223.
86. Abou-Kassem D, Kurita GP, Sjøgren P, Diasso PDK. Long-term opioid treatment and endocrine measures in patients with cancer-related pain: a systematic review. *Scand J Pain*. 2022;22(3):421-435.
87. Daniell HW. Hypogonadism in men consuming sustained-action oral opioids. *J Pain*. 2002;3(5):377-384.
88. Eshraghi Y, Hanks N, Rooney S, Ata FMY, Velasco C, Guirguis M, Uwaifo G. Establishing a Dose-Response Relationship Between Opioid Use and Hypogonadism: A Retrospective Case-Control Study. *Ochsner J*. 2021;21(3):249-253.
89. Rubinstein A, Carpenter DM. Elucidating risk factors for androgen deficiency associated with daily opioid use. *Am J Med*. 2014;127(12):1195-1201.
90. Rubinstein AL, Carpenter DM. Association Between Commonly Prescribed Opioids and Androgen Deficiency in Men: A Retrospective Cohort Analysis. *Pain Med*. 2017;18(4):637-644.
91. Bliesener N, Albrecht S, Schwager A, Weckbecker K, Lichtermann D, Klingmüller D. Plasma testosterone and sexual function in men receiving buprenorphine maintenance for opioid dependence. *J Clin Endocrinol Metab*. 2005;90(1):203-206.
92. Varma A, Sapra M, Iranmanesh A. Impact of opioid therapy on gonadal hormones: focus on buprenorphine. *Horm Mol Biol Clin Investig*. 2018;36(2).
93. Corsi DJ, Murphy MSQ. The Effects of opioids on female fertility, pregnancy and the breastfeeding mother-infant dyad: A Review. *Basic Clin Pharmacol Toxicol*. 2021;128(5):635-641.
94. Coluzzi F, Billeci D, Maggi M, Corona G. Testosterone deficiency in non-cancer opioid-treated patients. *J Endocrinol Invest*. 2018;41(12):1377-1388.
95. Bhasin S, Brito JP, Cunningham GR, Hayes FJ, Hodis HN, Matsumoto AM, Snyder PJ, Swerdloff RS, Wu FC, Yialamas MA. Testosterone Therapy in Men With Hypogonadism: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2018;103(5):1715-1744.
96. Fleseriu M, Hashim IA, Karavitaki N, Melmed S, Murad MH, Salvatori R, Samuels MH. Hormonal Replacement in Hypopituitarism in Adults: An Endocrine Society Clinical Practice Guideline. *J Clin Endocrinol Metab*. 2016;101(11):3888-3921.
97. Daniell HW, Lentz R, Mazer NA. Open-label pilot study of testosterone patch therapy in men with opioid-induced androgen deficiency. *J Pain*. 2006;7(3):200-210.
98. Aloisi AM, Ceccarelli I, Carlucci M, Suman A, Sindaco G, Mameli S, Paci V, Ravaioli L, Passavanti G, Bachiooco V, Pari G. Hormone replacement therapy in morphine-induced hypogonadic male chronic pain patients. *Reprod Biol Endocrinol*. 2011;9:26.
99. Basaria S, Travison TG, Alford D, Knapp PE, Teeter K, Cahalan C, Eder R, Lakshman K, Bachman E, Mensing G, Martel MO, Le D, Stroh H, Bhasin S, Wasan AD, Edwards RR. Effects of testosterone replacement in men with opioid-induced androgen deficiency: a randomized controlled trial. *Pain*. 2015;156(2):280-288.
100. Glintborg D, Vaegter HB, Christensen LL, Bendix E, Graven-Nielsen T, Andersen PG, Andersen M. Testosterone replacement therapy of opioid-induced male hypogonadism improved body composition but not pain perception: a double-blind, randomized, and placebo-controlled trial. *Eur J Endocrinol*. 2020;182(6):539-548.
101. Raheem OA, Patel SH, Sisul D, Furnish TJ, Hsieh TC. The Role of Testosterone Supplemental Therapy in Opioid-Induced Hypogonadism: A Retrospective Pilot Analysis. *Am J Mens Health*. 2017;11(4):1208-1213.
102. Kolind MI, Christensen LL, Caserotti P, Andersen MS, Glintborg D. Muscle function following testosterone replacement in men on opioid therapy for chronic non-cancer pain: A randomized controlled trial. *Andrology*. 2022;10(3):551-559.
103. Huang G, Travison T, Maggio M, Edwards RR, Basaria S. Effects of testosterone replacement on metabolic and inflammatory markers in men with opioid-induced androgen deficiency. *Clin Endocrinol (Oxf)*. 2016;85(2):232-238.

- 1637 104. Taylor T, Dluhy RG, Williams GH. beta-endorphin suppresses adrenocorticotropin and cortisol
1638 levels in normal human subjects. *J Clin Endocrinol Metab.* 1983;57(3):592-596.
- 1639 105. Stubbs WA, Delitala G, Jones A, Jeffcoate WJ, Edwards CR, Ratter SJ, Besser GM, Bloom
1640 SR, Alberti KG. Hormonal and metabolic responses to an enkephalin analogue in normal man.
1641 *Lancet.* 1978;2(8102):1225-1227.
- 1642 106. Rittmaster RS, Cutler GB, Jr., Sobel DO, Goldstein DS, Koppelman MC, Loriaux DL,
1643 Chrousos GP. Morphine inhibits the pituitary-adrenal response to ovine corticotropin-releasing
1644 hormone in normal subjects. *J Clin Endocrinol Metab.* 1985;60(5):891-895.
- 1645 107. Dackis CA, Gurpegui M, Pottash AL, Gold MS. Methadone induced hypoadrenalism. *Lancet.*
1646 1982;2(8308):1167.
- 1647 108. Facchinetti F, Volpe A, Farci G, Petraglia F, Porro CA, Barbieri G, Cioni A, Balestrieri A,
1648 Genazzani AR. Hypothalamus-pituitary-adrenal axis of heroin addicts. *Drug Alcohol Depend.*
1649 1985;15(4):361-366.
- 1650 109. Mendelson JH, Meyer RE, Ellingboe J, Mirin SM, McDougale M. Effects of heroin and
1651 methadone on plasma cortisol and testosterone. *J Pharmacol Exp Ther.* 1975;195(2):296-302.
- 1652 110. Palm S, Moenig H, Maier C. Effects of oral treatment with sustained release morphine tablets
1653 on hypothalamic-pituitary-adrenal axis. *Methods Find Exp Clin Pharmacol.* 1997;19(4):269-
1654 273.
- 1655 111. Müssig K, Knaus-Dittmann D, Schmidt H, Mörike K, Häring HU. Secondary adrenal failure
1656 and secondary amenorrhoea following hydromorphone treatment. *Clin Endocrinol (Oxf).*
1657 2007;66(4):604-605.
- 1658 112. Oltmanns KM, Fehm HL, Peters A. Chronic fentanyl application induces adrenocortical
1659 insufficiency. *J Intern Med.* 2005;257(5):478-480.
- 1660 113. Schimke KE, Greminger P, Brändle M. Secondary adrenal insufficiency due to opiate therapy
1661 - another differential diagnosis worth consideration. *Exp Clin Endocrinol Diabetes.*
1662 2009;117(10):649-651.
- 1663 114. Katz N, Mazer NA. The impact of opioids on the endocrine system. *Clin J Pain.*
1664 2009;25(2):170-175.
- 1665 115. Gibb FW, Stewart A, Walker BR, Strachan MW. Adrenal insufficiency in patients on long-
1666 term opioid analgesia. *Clin Endocrinol (Oxf).* 2016;85(6):831-835.
- 1667 116. Lamprecht A, Sorbello J, Jang C, Torpy DJ, Inder WJ. Secondary adrenal insufficiency and
1668 pituitary dysfunction in oral/transdermal opioid users with non-cancer pain. *Eur J Endocrinol.*
1669 2018;179(6):353-362.
- 1670 117. Li T, Cunningham JL, Gilliam WP, Loukianova L, Donegan DM, Bancos I. Prevalence of
1671 Opioid-Induced Adrenal Insufficiency in Patients Taking Chronic Opioids. *J Clin Endocrinol*
1672 *Metab.* 2020;105(10):e3766-3775.
- 1673 118. Rhodin A, Stridsberg M, Gordh T. Opioid endocrinopathy: a clinical problem in patients with
1674 chronic pain and long-term oral opioid treatment. *Clin J Pain.* 2010;26(5):374-380.
- 1675 119. Li T, Donegan D, Hooten WM, Bancos I. Clinical Presentation and Outcomes of Opioid-
1676 Induced Adrenal Insufficiency. *Endocr Pract.* 2020;26(11):1291-1297.
- 1677 120. Leow KP, Smith MT, Williams B, Cramond T. Single-dose and steady-state pharmacokinetics
1678 and pharmacodynamics of oxycodone in patients with cancer. *Clin Pharmacol Ther.*
1679 1992;52(5):487-495.
- 1680 121. Jannetto PJ, Bratanow NC. Utilization of pharmacogenomics and therapeutic drug monitoring
1681 for opioid pain management. *Pharmacogenomics.* 2009;10(7):1157-1167.
- 1682 122. Ali ES, Peiris AN. Secondary adrenal insufficiency induced by long-term use of opioid
1683 analgesics. *Proc (Bayl Univ Med Cent).* 2019;32(3):417-418.
- 1684 123. Lee AS, Twigg SM. Opioid-induced secondary adrenal insufficiency presenting as
1685 hypercalcaemia. *Endocrinol Diabetes Metab Case Rep.* 2015;2015:150035.
- 1686 124. Tabet EJ, Clarke AJ, Twigg SM. Opioid-induced hypoadrenalism resulting in fasting
1687 hypoglycaemia. *BMJ Case Rep.* 2019;12(12).
- 1688 125. Werumeus Buning J, Brummelman P, Koerts J, Dullaart RP, van den Berg G, van der Klauw
1689 MM, Sluiter WJ, Tucha O, Wolffenbuttel BH, van Beek AP. Hydrocortisone Dose Influences
1690 Pain, Depressive Symptoms and Perceived Health in Adrenal Insufficiency: A Randomized
1691 Controlled Trial. *Neuroendocrinology.* 2016;103(6):771-778.

- 1692 126. Husebye ES, Pearce SH, Krone NP, Kämpe O. Adrenal insufficiency. *Lancet*.
1693 2021;397(10274):613-629.
- 1694 127. Al-Aridi R, Abdelmannan D, Arafah BM. Biochemical diagnosis of adrenal insufficiency: the
1695 added value of dehydroepiandrosterone sulfate measurements. *Endocr Pract*. 2011;17(2):261-
1696 270.
- 1697 128. Sarlos S, Inder WJ. Selective use of the insulin tolerance test to diagnose hypopituitarism.
1698 *Intern Med J*. 2013;43(1):89-93.
- 1699 129. Soule S, Van Zyl Smit C, Parolis G, Attenborough S, Peter D, Kinvig S, Kinvig T, Coetzer E.
1700 The low dose ACTH stimulation test is less sensitive than the overnight metyrapone test for the
1701 diagnosis of secondary hypoadrenalism. *Clin Endocrinol (Oxf)*. 2000;53(2):221-227.
- 1702 130. English KA, Chikani V, Jang C, Dimeski G, Olson S, Inder WJ. The relationship between early
1703 post-operative ACTH / cortisol following pituitary surgery and long-term glucocorticoid
1704 requirement - Do ultradian rhythms matter? *Clin Endocrinol (Oxf)*. 2021;94(4):636-644.
- 1705 131. Colling C, Nachtigall L, Biller BMK, Miller KK. The biochemical diagnosis of adrenal
1706 insufficiency with modern cortisol assays: Reappraisal in the setting of opioid exposure and
1707 hospitalization. *Clin Endocrinol (Oxf)*. 2022;96(1):21-29.
- 1708 132. Nenke MA, Haylock CL, Rankin W, Inder WJ, Gagliardi L, Eldridge C, Rolan P, Torpy DJ.
1709 Low-dose hydrocortisone replacement improves wellbeing and pain tolerance in chronic pain
1710 patients with opioid-induced hypocortisolemic responses. A pilot randomized, placebo-
1711 controlled trial. *Psychoneuroendocrinology*. 2015;56:157-167.
- 1712 133. Bhansali A, Velayutham P, Sialy R, Sethi B. Effect of opiates on growth hormone secretion in
1713 acromegaly. *Horm Metab Res*. 2005;37(7):425-427.
- 1714 134. Wehrenberg WB, Bloch B, Ling N. Pituitary secretion of growth hormone in response to opioid
1715 peptides and opiates is mediated through growth hormone-releasing factor.
1716 *Neuroendocrinology*. 1985;41(1):13-16.
- 1717 135. Delitala G, Tomasi PA, Palermo M, Ross RJ, Grossman A, Besser GM. Opioids stimulate
1718 growth hormone (GH) release in man independently of GH-releasing hormone. *J Clin*
1719 *Endocrinol Metab*. 1989;69(2):356-358.
- 1720 136. Valverde-Filho J, da Cunha Neto MB, Fonoff ET, Meirelles Ede S, Teixeira MJ. Chronic spinal
1721 and oral morphine-induced neuroendocrine and metabolic changes in noncancer pain patients.
1722 *Pain Med*. 2015;16(4):715-725.
- 1723 137. Merza Z, Edwards N, Walters SJ, Newell-Price J, Ross RJ. Patients with chronic pain and
1724 abnormal pituitary function require investigation. *Lancet*. 2003;361(9376):2203-2204.
- 1725 138. Cushman P, Jr. Growth hormone in narcotic addiction. *J Clin Endocrinol Metab*.
1726 1972;35(3):352-358.
- 1727 139. Sustkova-Fiserova M, Charalambous C, Khryakova A, Certilina A, Lapka M, Šlamberová R.
1728 The Role of Ghrelin/GHS-R1A Signaling in Nonalcohol Drug Addictions. *Int J Mol Sci*.
1729 2022;23(2).
- 1730 140. Rhodin A, von Ehren M, Skottheim B, Grönbladh A, Ortiz-Nieto F, Raininko R, Gordh T,
1731 Nyberg F. Recombinant human growth hormone improves cognitive capacity in a pain patient
1732 exposed to chronic opioids. *Acta Anaesthesiol Scand*. 2014;58(6):759-765.
- 1733 141. The Pituitary. Drugs and Pituitary Function. Monica R. Gadelha 5th ed: Academic Press.
- 1734 142. Tolis G, Dent R, Guyda H. Opiates, prolactin, and the dopamine receptor. *J Clin Endocrinol*
1735 *Metab*. 1978;47(1):200-203.
- 1736 143. Hoehe M, Duka T, Doenicke A. Human studies on the mu opiate receptor agonist fentanyl:
1737 neuroendocrine and behavioral responses. *Psychoneuroendocrinology*. 1988;13(5):397-408.
- 1738 144. Mendelson JH, Mello NK, Teoh SK, Lloyd-Jones JG, Clifford JM. Naloxone suppresses
1739 buprenorphine stimulation of plasma prolactin. *J Clin Psychopharmacol*. 1989;9(2):105-109.
- 1740 145. Saarialho-Kere U, Mattila MJ, Seppälä T. Psychomotor, respiratory and neuroendocrinological
1741 effects of a mu-opioid receptor agonist (oxycodone) in healthy volunteers. *Pharmacol Toxicol*.
1742 1989;65(4):252-257.
- 1743 146. Zis AP, Haskett RF, Albala AA, Carroll BJ. Morphine inhibits cortisol and stimulates prolactin
1744 secretion in man. *Psychoneuroendocrinology*. 1984;9(4):423-427.

- 1745 147. Duarte RV, Raphael JH, Southall JL, Labib MH, Whallett AJ, Ashford RL. Hypogonadism and
1746 low bone mineral density in patients on long-term intrathecal opioid delivery therapy. *BMJ*
1747 *Open*. 2013;3(6).
- 1748 148. Wong D, Gray DP, Simmonds M, Rashid S, Sobolev I, Morrish DW. Opioid analgesics
1749 suppress male gonadal function but opioid use in males and females does not correlate with
1750 symptoms of sexual dysfunction. *Pain Res Manag*. 2011;16(5):311-316.
- 1751 149. Moshtaghi-Kashanian GR, Esmaeili F, Dabiri S. Enhanced prolactin levels in opium smokers.
1752 *Addict Biol*. 2005;10(4):345-349.
- 1753 150. Kley HK, Oellerich M, Wiegelmann W, Herrmann J, Rudorff KH, Nieschlag E, Krüskemper
1754 HL. The effect of methadone on hypophyseal and peripheral glandular hormones during
1755 withdrawal. *Horm Metab Res*. 1977;9(6):484-488.
- 1756 151. Vescovi PP, Pezzarossa A, Ceresini G, Rastelli G, Valenti G, Gerra G. Effects of dopamine
1757 receptor stimulation on opiate-induced modifications of pituitary-gonadal function. *Horm Res*.
1758 1985;21(3):155-159.
- 1759 152. Petersenn S, Fleseriu M, Casanueva FF, Giustina A, Biermasz N, Biller BMK, Bronstein M,
1760 Chanson P, Fukuoka H, Gadelha M, Greenman Y, Gurnell M, Ho KKY, Honegger J,
1761 Ioachimescu AG, Kaiser UB, Karavitaki N, Katznelson L, Lodish M, Maiter D, Marcus HJ,
1762 McCormack A, Molitch M, Muir CA, Neggers S, Pereira AM, Pivonello R, Post K, Raverot G,
1763 Salvatori R, Samson SL, Shimon I, Spencer-Segal J, Vila G, Wass J, Melmed S. Diagnosis and
1764 management of prolactin-secreting pituitary adenomas: a Pituitary Society international
1765 Consensus Statement. *Nat Rev Endocrinol*. 2023;19(12):722-740.
- 1766 153. Devilla L, Pende A, Morgano A, Giusti M, Musso NR, Lotti G. Morphine-induced TSH release
1767 in normal and hypothyroid subjects. *Neuroendocrinology*. 1985;40(4):303-308.
- 1768 154. Pende A, Musso NR, Montaldi ML, Pastorino G, Arzese M, Devilla L. Evaluation of the effects
1769 induced by four opiate drugs, with different affinities to opioid receptor subtypes, on anterior
1770 pituitary LH, TSH, PRL and GH secretion and on cortisol secretion in normal men. *Biomed*
1771 *Pharmacother*. 1986;40(5):178-182.
- 1772 155. Brambilla F, Nobile P, Zanoboni A, Zanoboni-Muciaccia W, Meroni PL. Effects of chronic
1773 heroin addiction on pituitary-thyroid function in man. *J Endocrinol Invest*. 1980;3(3):251-255.
- 1774 156. Vescovi PP, Gerra G, Rastelli G, Ceda GP, Valenti G. Effect of methadone on TSH and thyroid
1775 hormone secretion. *Horm Metab Res*. 1984;16(1):53-54.
- 1776 157. Spagnoli W, De Venuto G, Mattarei M, Dal Ri P, Miori R. Prolactin and thyrotropin pituitary
1777 response to thyrotropin releasing hormone in young female heroin addicts. *Drug Alcohol*
1778 *Depend*. 1987;20(3):247-254.
- 1779 158. Fraser LA, Morrison D, Morley-Forster P, Paul TL, Tokmakejian S, Larry Nicholson R, Bureau
1780 Y, Friedman TC, Van Uum SH. Oral opioids for chronic non-cancer pain: higher prevalence of
1781 hypogonadism in men than in women. *Exp Clin Endocrinol Diabetes*. 2009;117(1):38-43.
- 1782 159. Gozashti MH, Mohammadzadeh E, Divsalar K, Shokoohi M. The effect of opium addiction on
1783 thyroid function tests. *J Diabetes Metab Disord*. 2014;13(1):5.
- 1784 160. Shenkman L, Massie B, Mitsuma T, Hollander CS. Effects of chronic methadone
1785 administration on the hypothalamic-pituitary-thyroid axis. *J Clin Endocrinol Metab*.
1786 1972;35(1):169-170.
- 1787 161. Pezeshki B, Pourmontaseri H, Homayounfar R, Moghaddam MT, Dehghan A. Thyroid function
1788 and opium use disorder: a cross-sectional study on the Fasa adults cohort study (FACS), 2017.
1789 *BMC Endocr Disord*. 2023;23(1):264.
- 1790 162. Pan Y, Xu R. Mining comorbidities of opioid use disorder from FDA adverse event reporting
1791 system and patient electronic health records. *BMC Med Inform Decis Mak*. 2022;22(Suppl
1792 2):155.
- 1793 163. R. C. De B. THE ANTIDIURETIC ACTION OF MORPHINE, AND ITS MECHANISM.
1794 *Journal of Pharmacology and Experimental Therapeutics*. 1944;82(1):74.
- 1795 164. Schnieden H, Blackmore EK. The effect of nalorphine on the antidiuretic action of morphine
1796 in rats and men. *Br J Pharmacol Chemother*. 1955;10(1):45-50.
- 1797 165. Bozkurt P, Kaya G, Yeker Y, Altintas F, Bakan M, Hacibekiroglu M, Kavunoglu G. Effects of
1798 systemic and epidural morphine on antidiuretic hormone levels in children. *Paediatr Anaesth*.
1799 2003;13(6):508-514.

- 1800 166. Boulton AJ, Wilson N, Turnbull KW, Yip RW. Haemodynamic and plasma vasopressin
1801 responses during high-dose fentanyl or sufentanil anaesthesia. *Can Anaesth Soc J*.
1802 1986;33(4):475-483.
- 1803 167. Weiskopf RB, Reid IA, Fisher DM, Holmes MA, Rosen JI, Keil LC. Effects of fentanyl on
1804 vasopressin secretion in human subjects. *J Pharmacol Exp Ther*. 1987;242(3):970-973.
- 1805 168. Liamis G, Milionis H, Elisaf M. A review of drug-induced hyponatremia. *Am J Kidney Dis*.
1806 2008;52(1):144-153.
- 1807 169. Langfeldt LA, Cooley ME. Syndrome of inappropriate antidiuretic hormone secretion in
1808 malignancy: review and implications for nursing management. *Clin J Oncol Nurs*.
1809 2003;7(4):425-430.
- 1810 170. Hunter R. Tramadol and hyponatraemia. *Australian Prescriber*. 2004:27-97.
- 1811 171. Abadie D, Durrieu G, Roussin A, Montastruc JL. ["Serious" adverse drug reactions with
1812 tramadol: a 2010-2011 pharmacovigilance survey in France]. *Therapie*. 2013;68(2):77-84.
- 1813 172. Le Berre JP, Desramé J, Lecoules S, Coutant G, Béchade D, Algayres JP. [Hyponatremia due
1814 to tramadol]. *Rev Med Interne*. 2007;28(12):888-889.
- 1815 173. Lota AS, Dubrey SW, Wills P. Profound hyponatraemia following a tramadol overdose. *Qjm*.
1816 2012;105(4):397-398.
- 1817 174. Fournier JP, Yin H, Nessim SJ, Montastruc JL, Azoulay L. Tramadol for noncancer pain and
1818 the risk of hyponatremia. *Am J Med*. 2015;128(4):418-425.e415.
- 1819 175. American Geriatrics Society 2019 Updated AGS Beers Criteria® for Potentially Inappropriate
1820 Medication Use in Older Adults. *J Am Geriatr Soc*. 2019;67(4):674-694.
- 1821 176. Virnes RE, Tiihonen M, Karttunen N, van Poelgeest EP, van der Velde N, Hartikainen S.
1822 Opioids and Falls Risk in Older Adults: A Narrative Review. *Drugs Aging*. 2022;39(3):199-
1823 207.
- 1824 177. Grossman A, Besser GM, Milles JJ, Baylis PH. Inhibition of vasopressin release in man by an
1825 opiate peptide. *Lancet*. 1980;2(8204):1108-1110.
- 1826 178. Brownell J, del Pozo E, Donatsch P. Inhibition of vasopressin secretion by a met-enkephalin
1827 (FK 33-824) in humans. *Acta Endocrinol (Copenh)*. 1980;94(3):304-308.
- 1828 179. Lightman SL, Forsling ML. The effect of the methionine enkephalin analogue DAMME on the
1829 vasopression response to tilt in man. *Clin Sci (Lond)*. 1980;59(6):501-503.
- 1830 180. Kramer HJ, Uhl W, Ladstetter B, Bäcker A. Influence of asimadoline, a new kappa-opioid
1831 receptor agonist, on tubular water absorption and vasopressin secretion in man. *Br J Clin*
1832 *Pharmacol*. 2000;50(3):227-235.
- 1833 181. Glahn A, Heberlein A, Dürsteler-MacFarland KM, Lenz B, Frieling H, Gröschl M, Wiesbeck
1834 GA, Kornhuber J, Bönsch D, Bleich S, Hillemacher T. Atrial natriuretic peptide, arginine
1835 vasopressin peptide and cortisol serum levels in opiate-dependent patients.
1836 *Neuropsychobiology*. 2013;67(2):111-115.
- 1837 182. Bhat N, Balliu E, Osipoff J, Lane A, Wilson T. Opioid-induced hyponatremia in a patient with
1838 central diabetes insipidus: independence from ADH. *J Pediatr Endocrinol Metab*.
1839 2017;30(6):693-696.
- 1840 183. Land H, Schütz G, Schmale H, Richter D. Nucleotide sequence of cloned cDNA encoding
1841 bovine arginine vasopressin-neurophysin II precursor. *Nature*. 1982;295(5847):299-303.
- 1842 184. Balanescu S, Kopp P, Gaskill MB, Morgenthaler NG, Schindler C, Rutishauser J. Correlation
1843 of plasma copeptin and vasopressin concentrations in hypo-, iso-, and hyperosmolar States. *J*
1844 *Clin Endocrinol Metab*. 2011;96(4):1046-1052.
- 1845 185. Blum CA, Velly L, Brochet C, Ziegler F, Tavolacci MP, Hausfater P, Lvovschi VE. Relevance
1846 of cortisol and copeptin blood concentration changes in an experimental pain model. *Sci Rep*.
1847 2022;12(1):4767.
- 1848 186. Domes G, Heinrichs M, Gläscher J, Büchel C, Braus DF, Herpertz SC. Oxytocin attenuates
1849 amygdala responses to emotional faces regardless of valence. *Biol Psychiatry*.
1850 2007;62(10):1187-1190.
- 1851 187. Heinrichs M, von Dawans B, Domes G. Oxytocin, vasopressin, and human social behavior.
1852 *Front Neuroendocrinol*. 2009;30(4):548-557.

- 1853 188. Meyer-Lindenberg A, Domes G, Kirsch P, Heinrichs M. Oxytocin and vasopressin in the
1854 human brain: social neuropeptides for translational medicine. *Nat Rev Neurosci*.
1855 2011;12(9):524-538.
- 1856 189. Neumann ID, Slattery DA. Oxytocin in General Anxiety and Social Fear: A Translational
1857 Approach. *Biol Psychiatry*. 2016;79(3):213-221.
- 1858 190. Heinrichs M, Baumgartner T, Kirschbaum C, Ehler U. Social support and oxytocin interact to
1859 suppress cortisol and subjective responses to psychosocial stress. *Biol Psychiatry*.
1860 2003;54(12):1389-1398.
- 1861 191. Ludwig M, Leng G. Dendritic peptide release and peptide-dependent behaviours. *Nat Rev*
1862 *Neurosci*. 2006;7(2):126-136.
- 1863 192. Kirsch P, Esslinger C, Chen Q, Mier D, Lis S, Siddhanti S, Gruppe H, Mattay VS, Gallhofer
1864 B, Meyer-Lindenberg A. Oxytocin modulates neural circuitry for social cognition and fear in
1865 humans. *J Neurosci*. 2005;25(49):11489-11493.
- 1866 193. Lindow SW, van der Spuy ZM, Hendricks MS, Nugent FA, Dunne TT. The effect of morphine
1867 and naloxone administration on maternal oxytocin concentration in late pregnancy. *Clin*
1868 *Endocrinol (Oxf)*. 1993;39(6):671-675.
- 1869 194. Lindow SW, Hendricks MS, Thompson JW, van der Spuy ZM. Effects of morphine
1870 administration on the fetal production of oxytocin in labour. *Clin Sci (Lond)*. 1998;95(1):91-
1871 95.
- 1872 195. Lindow SW, van der Spuy ZM, Hendricks MS, Rosselli AP, Lombard C, Leng G. The effect
1873 of morphine and naloxone administration on plasma oxytocin concentrations in the first stage
1874 of labour. *Clin Endocrinol (Oxf)*. 1992;37(4):349-353.
- 1875 196. Lindow SW, Hendricks MS, Nugent FA, Dunne TT, van der Spuy ZM. Morphine suppresses
1876 the oxytocin response in breast-feeding women. *Gynecol Obstet Invest*. 1999;48(1):33-37.
- 1877 197. Shibli KU, Dhillon AR, Goode JA, Gilbert CL, Thompson JW, Russell IF, Lindow SW. Effect
1878 of intrathecal fentanyl on oxytocin secretion in pregnant women not in labour. *Clin Sci (Lond)*.
1879 2001;101(4):415-419.
- 1880 198. Stocche RM, Klamt JG, Antunes-Rodrigues J, Garcia LV, Moreira AC. Effects of intrathecal
1881 sufentanil on plasma oxytocin and cortisol concentrations in women during the first stage of
1882 labor. *Reg Anesth Pain Med*. 2001;26(6):545-550.
- 1883 199. Miyakoshi K, Tanaka M, Morisaki H, Kim SH, Hosokawa Y, Matsumoto T, Minegishi K,
1884 Yoshimura Y. Perinatal outcomes: intravenous patient-controlled fentanyl versus no analgesia
1885 in labor. *J Obstet Gynaecol Res*. 2013;39(4):783-789.
- 1886 200. Anim-Somuah M, Smyth RM, Cyna AM, Cuthbert A. Epidural versus non-epidural or no
1887 analgesia for pain management in labour. *Cochrane Database Syst Rev*. 2018;5(5):Cd000331.
- 1888 201. Moeini M, Omid A, Sehat M, Banafshe HR. The Effects of Oxytocin on Withdrawal, Craving
1889 and Stress Response in Heroin-Dependent Patients: A Randomized, Double-Blind Clinical
1890 Trial. *Eur Addict Res*. 2019;25(1):41-47.
- 1891 202. Buisman-Pijlman FT, Sumracki NM, Gordon JJ, Hull PR, Carter CS, Tops M. Individual
1892 differences underlying susceptibility to addiction: Role for the endogenous oxytocin system.
1893 *Pharmacol Biochem Behav*. 2014;119:22-38.
- 1894 203. Tops M, Koole SL, H IJ, Buisman-Pijlman FT. Why social attachment and oxytocin protect
1895 against addiction and stress: Insights from the dynamics between ventral and dorsal
1896 corticostriatal systems. *Pharmacol Biochem Behav*. 2014;119:39-48.
- 1897 204. Coluzzi F, Pergolizzi J, Raffa RB, Mattia C. The unsolved case of "bone-impairing analgesics":
1898 the endocrine effects of opioids on bone metabolism. *Ther Clin Risk Manag*. 2015;11:515-523.
- 1899 205. Carvalho AL, Brooks DJ, Barlow D, Langlais AL, Morrill B, Houseknecht KL, Bouxsein ML,
1900 Lian JB, King T, Farina NH, Motyl KJ. Sustained Morphine Delivery Suppresses Bone
1901 Formation and Alters Metabolic and Circulating miRNA Profiles in Male C57BL/6J Mice. *J*
1902 *Bone Miner Res*. 2022;37(11):2226-2243.
- 1903 206. Holan V, Cechova K, Zajicova A, Koss J, Hermankova B, Bohacova P, Hajkova M, Krulova
1904 M, Svoboda P, Javorkova E. The Impact of Morphine on the Characteristics and Function
1905 Properties of Human Mesenchymal Stem Cells. *Stem Cell Rev Rep*. 2018;14(6):801-811.
- 1906 207. Marczak ED, Jinsmaa Y, Myers PH, Blankenship T, Wilson R, Balboni G, Salvadori S, Lazarus
1907 LH. Orally administered H-Dmt-Tic-Lys-NH-CH₂-Ph (MZ-2), a potent mu/delta-opioid

- 1908 receptor antagonist, regulates obese-related factors in mice. *Eur J Pharmacol.* 2009;616(1-
1909 3):115-121.
- 1910 208. Pérez-Castrillón JL, Olmos JM, Gómez JJ, Barrallo A, Riancho JA, Perera L, Valero C, Amado
1911 JA, González-Macías J. Expression of opioid receptors in osteoblast-like MG-63 cells, and
1912 effects of different opioid agonists on alkaline phosphatase and osteocalcin secretion by these
1913 cells. *Neuroendocrinology.* 2000;72(3):187-194.
- 1914 209. Saeidinezhad M, Razban V, Safizadeh H, Ezzatabadipour M. Effects of maternal consumption
1915 of morphine on rat skeletal system development. *BMC Musculoskelet Disord.* 2021;22(1):435.
- 1916 210. Tanaka K, Kondo H, Hamamura K, Togari A. Systemic administration of low-dose naltrexone
1917 increases bone mass due to blockade of opioid growth factor receptor signaling in mice
1918 osteoblasts. *Life Sci.* 2019;224:232-240.
- 1919 211. Hosseini-Fard SR, Etemad-Moghadam S, Alaeddini M, Dehpour AR, Emamgholipour S,
1920 Golestani A. Exploring the impact of naltrexone on the THBS1/eNOS/NO pathway in
1921 osteoporotic bile duct-ligated rats. *Sci Rep.* 2024;14(1):48.
- 1922 212. Moradi M, Doustimotlagh AH, Dehpour AR, Rahimi N, Golestani A. The influence of TRAIL,
1923 adiponectin and sclerostin alterations on bone loss in BDL-induced cirrhotic rats and the effect
1924 of opioid system blockade. *Life Sci.* 2019;233:116706.
- 1925 213. Rico H, Costales C, Cabranes JA, Escudero M. Lower serum osteocalcin levels in pregnant
1926 drug users and their newborns at the time of delivery. *Obstet Gynecol.* 1990;75(6):998-1000.
- 1927 214. Boshra V. Evaluation of osteoporosis risk associated with chronic use of morphine, fentanyl
1928 and tramadol in adult female rats. *Curr Drug Saf.* 2011;6(3):159-163.
- 1929 215. Wilhelm W, Kreuer S. The place for short-acting opioids: special emphasis on remifentanyl.
1930 *Crit Care.* 2008;12 Suppl 3(Suppl 3):S5.
- 1931 216. Baik SW, Park BS, Kim YH, Kim YD, Kim CH, Yoon JY, Yoon JU. Effects of Remifentanyl
1932 Preconditioning on Osteoblasts under Hypoxia-Reoxygenation Condition. *Int J Med Sci.*
1933 2015;12(7):583-589.
- 1934 217. Yoon JY, Kim TS, Ahn JH, Yoon JU, Kim HJ, Kim EJ. Remifentanyl promotes
1935 osteoblastogenesis by upregulating Runx2/osterix expression in preosteoblastic C2C12 cells. *J*
1936 *Dent Anesth Pain Med.* 2019;19(2):91-99.
- 1937 218. Hioki T, Tokuda H, Tanabe K, Kim W, Tachi J, Yamaguchi S, Matsushima-Nishiwaki R,
1938 Kozawa O, Iida H. Amplification by tramadol of PGD(2)-induced osteoprotegerin synthesis in
1939 osteoblasts: Involvement of μ -opioid receptor and 5-HT transporter. *Prostaglandins Leukot*
1940 *Essent Fatty Acids.* 2021;172:102323.
- 1941 219. Kim W, Tanabe K, Kuroyanagi G, Matsushima-Nishiwaki R, Fujita K, Kawabata T, Sakai G,
1942 Tachi J, Hioki T, Nakashima D, Yamaguchi S, Otsuka T, Tokuda H, Kozawa O, Iida H.
1943 Tramadol enhances PGF(2 α)-stimulated osteoprotegerin synthesis in osteoblasts. *Heliyon.*
1944 2020;6(8):e04779.
- 1945 220. Jeon HO, Choi IS, Yoon JY, Kim EJ, Yoon JU, Cho AR, Kim HJ, Kim CH. Effect of
1946 remifentanyl on pre-osteoclast cell differentiation in vitro. *J Dent Anesth Pain Med.*
1947 2018;18(1):9-17.
- 1948 221. Yoon JY, Baek CW, Kim HJ, Kim EJ, Byeon GJ, Yoon JU. Remifentanyl Negatively Regulates
1949 RANKL-Induced Osteoclast Differentiation and Bone Resorption by Inhibiting c-Fos/NFATc1
1950 Expression. *Tissue Eng Regen Med.* 2018;15(3):333-340.
- 1951 222. Balodimos S, Nikolaou K, Njau S, Karamouzis M, Kovatsi L. The effect of opioid dependence
1952 on conventional and novel biochemical parameters of bone metabolism. *Am J Drug Alcohol*
1953 *Abuse.* 2015;41(6):535-540.
- 1954 223. Gotthardt F, Huber C, Thierfelder C, Grize L, Kraenzlin M, Scheidegger C, Meier C. Bone
1955 mineral density and its determinants in men with opioid dependence. *J Bone Miner Metab.*
1956 2017;35(1):99-107.
- 1957 224. Grey A, Rix-Trott K, Horne A, Gamble G, Bolland M, Reid IR. Decreased bone density in men
1958 on methadone maintenance therapy. *Addiction.* 2011;106(2):349-354.
- 1959 225. Coluzzi F, Scerpa MS, Centanni M. The Effect of Opiates on Bone Formation and Bone
1960 Healing. *Curr Osteoporos Rep.* 2020;18(3):325-335.
- 1961 226. Chrastil J, Sampson C, Jones KB, Higgins TF. Postoperative opioid administration inhibits
1962 bone healing in an animal model. *Clin Orthop Relat Res.* 2013;471(12):4076-4081.

- 1963 227. Jain N, Himed K, Toth JM, Briley KC, Phillips FM, Khan SN. Opioids delay healing of spinal
1964 fusion: a rabbit posterolateral lumbar fusion model. *Spine J.* 2018;18(9):1659-1668.
- 1965 228. Petrizzi L, Mariscoli M, Valbonetti L, Varasano V, Langhoff JD, Von Rechenberg B.
1966 Preliminary study on the effect of parenteral naloxone, alone and in association with calcium
1967 gluconate, on bone healing in an ovine "drill hole" model system. *BMC Musculoskelet Disord.*
1968 2007;8:43.
- 1969 229. Kaye AD, Baluch A, Scott JT. Pain management in the elderly population: a review. *Ochsner*
1970 *J.* 2010;10(3):179-187.
- 1971 230. Chau DL, Walker V, Pai L, Cho LM. Opiates and elderly: use and side effects. *Clin Interv*
1972 *Aging.* 2008;3(2):273-278.
- 1973 231. Al-Qurain AA, Gebremichael LG, Khan MS, Williams DB, Mackenzie L, Phillips C, Russell
1974 P, Roberts MS, Wiese MD. Prevalence and Factors Associated with Analgesic Prescribing in
1975 Poly-Medicated Elderly Patients. *Drugs Aging.* 2020;37(4):291-300.
- 1976 232. Mörttinen-Vallius H, Hartikainen S, Seinela L, Jämsen E. The prevalence of and exact
1977 indications for daily opioid use among aged home care clients with and without dementia. *Aging*
1978 *Clin Exp Res.* 2021;33(5):1239-1247.
- 1979 233. Torchia MT, Munson J, Tosteson TD, Tosteson ANA, Wang Q, McDonough CM, Morgan TS,
1980 Bynum JPW, Bell JE. Patterns of Opioid Use in the 12 Months Following Geriatric Fragility
1981 Fractures: A Population-Based Cohort Study. *J Am Med Dir Assoc.* 2019;20(3):298-304.
- 1982 234. Zullo AR, Zhang T, Beaudoin FL, Lee Y, McConeghy KW, Kiel DP, Daiello LA, Mor V, Berry
1983 SD. Pain Treatments After Hip Fracture Among Older Nursing Home Residents. *J Am Med Dir*
1984 *Assoc.* 2018;19(2):174-176.
- 1985 235. Ni W, Ricker C, Quinn M, Gasquet N, Janardhanan D, Gilligan CJ, Hirsch JA. Trends in opioid
1986 use following balloon kyphoplasty or vertebroplasty for the treatment of vertebral compression
1987 fractures. *Osteoporos Int.* 2022;33(4):821-837.
- 1988 236. Tolba R, Bolash RB, Shroll J, Costandi S, Dalton JE, Sanghvi C, Mekhail N. Kyphoplasty
1989 increases vertebral height, decreases both pain score and opiate requirements while improving
1990 functional status. *Pain Pract.* 2014;14(3):E91-97.
- 1991 237. Yoshida K, Yu Z, Greendale GA, Ruppert K, Lian Y, Tedeschi SK, Lin TC, Haneuse S, Glynn
1992 RJ, Hernández-Díaz S, Solomon DH. Effects of analgesics on bone mineral density: A
1993 longitudinal analysis of the prospective SWAN cohort with three-group matching weights.
1994 *Pharmacoepidemiol Drug Saf.* 2018;27(2):182-190.
- 1995 238. Vestergaard P, Hermann P, Jensen JE, Eiken P, Mosekilde L. Effects of paracetamol, non-
1996 steroidal anti-inflammatory drugs, acetylsalicylic acid, and opioids on bone mineral density and
1997 risk of fracture: results of the Danish Osteoporosis Prevention Study (DOPS). *Osteoporos Int.*
1998 2012;23(4):1255-1265.
- 1999 239. Finch PM, Price LM, Pullan PT, Drummond PD. Effects of testosterone treatment on bone
2000 mineral density in hypogonadal men receiving intrathecal opioids. *Pain Pract.* 2015;15(4):308-
2001 313.
- 2002 240. Kinjo M, Setoguchi S, Schneeweiss S, Solomon DH. Bone mineral density in subjects using
2003 central nervous system-active medications. *Am J Med.* 2005;118(12):1414.
- 2004 241. Milos G, Gallo LM, Sosic B, Uebelhart D, Goerres G, Haeuselmann HJ, Eich D. Bone mineral
2005 density in young women on methadone substitution. *Calcif Tissue Int.* 2011;89(3):228-233.
- 2006 242. Kim TW, Alford DP, Malabanan A, Holick MF, Samet JH. Low bone density in patients
2007 receiving methadone maintenance treatment. *Drug Alcohol Depend.* 2006;85(3):258-262.
- 2008 243. Sharma A, Cohen HW, Freeman R, Santoro N, Schoenbaum EE. Prospective evaluation of
2009 bone mineral density among middle-aged HIV-infected and uninfected women: Association
2010 between methadone use and bone loss. *Maturitas.* 2011;70(3):295-301.
- 2011 244. Ramli FF, Syed Hashim SA, Mohd Effendy N. Factors Associated with Low Bone Density in
2012 Opioid Substitution Therapy Patients: A Systematic Review. *Int J Med Sci.* 2021;18(2):575-
2013 581.
- 2014 245. Ping F, Wang Y, Wang J, Chen J, Zhang W, Zhi H, Liu Y. Opioids increase hip fracture risk:
2015 a meta-analysis. *J Bone Miner Metab.* 2017;35(3):289-297.
- 2016 246. Teng Z, Zhu Y, Wu F, Zhu Y, Zhang X, Zhang C, Wang S, Zhang L. Opioids contribute to
2017 fracture risk: a meta-analysis of 8 cohort studies. *PLoS One.* 2015;10(6):e0128232.

- 2018 247. Yoshikawa A, Ramirez G, Smith ML, Foster M, Nabil AK, Jani SN, Ory MG. Opioid Use and
2019 the Risk of Falls, Fall Injuries and Fractures among Older Adults: A Systematic Review and
2020 Meta-Analysis. *J Gerontol A Biol Sci Med Sci*. 2020;75(10):1989-1995.
- 2021 248. Yue Q, Ma Y, Teng Y, Zhu Y, Liu H, Xu S, Liu J, Liu J, Zhang X, Teng Z. An updated analysis
2022 of opioids increasing the risk of fractures. *PLoS One*. 2020;15(4):e0220216.
- 2023 249. Guan Q, Men S, Juurlink DN, Bronskill SE, Wunsch H, Gomes T. Opioid Initiation and the
2024 Hazard of Falls or Fractures Among Older Adults with Varying Levels of Central Nervous
2025 System Depressant Burden. *Drugs Aging*. 2022;39(9):729-738.
- 2026 250. Taipale H, Hamina A, Karttunen N, Koponen M, Tanskanen A, Tiihonen J, Hartikainen S,
2027 Tolppanen AM. Incident opioid use and risk of hip fracture among persons with Alzheimer
2028 disease: a nationwide matched cohort study. *Pain*. 2019;160(2):417-423.
- 2029 251. Silverman S, Schepman P, Rice JB, Beck CG, Pajeroski W, White AG, Thakkar S, Robinson
2030 RL, Emir B. Risk Factors Associated with Falls and Fractures Following Prescription of
2031 Opioids Among Privately Insured Patients with Osteoarthritis. *J Health Econ Outcomes Res*.
2032 2022;9(2):47-56.
- 2033 252. Tveit M, Rosengren BE, Nilsson JA, Karlsson MK. Exercise in youth: High bone mass, large
2034 bone size, and low fracture risk in old age. *Scand J Med Sci Sports*. 2015;25(4):453-461.
- 2035 253. Ding Z, Ma Z, Yang X, Sun Y. Effect of Eight-Month Exercise Intervention on Bone Outcomes
2036 of Young Opioid-Dependent Women. *Int J Environ Res Public Health*. 2021;18(21).

2037