



FROM THE EXPERTS IN ENDOCRINOLOGY

2026 ENDOCRINE CASE MANAGEMENT: MEET THE PROFESSOR

FROM THE EXPERTS IN ENDOCRINOLOGY

ENDO 2026

MEET THE PROFESSOR



ENDOCRINE

CASE MANAGEMENT

ENDO  2026

ENDOCRINE
SOCIETY 



2055 L Street, NW, Suite 600
Washington, DC 20036
www.endocrine.org

Other Publications:
press.endocrine.org

For between-edition updates, visit us at:
endocrine.org/education-and-training/book-updates

The Endocrine Society is the world's largest, oldest, and most active organization working to advance the clinical practice of endocrinology and hormone research. Founded in 1916, the Society now has more than 18,000 global members across a range of disciplines.

The Society has earned an international reputation for excellence in the quality of its peer-reviewed journals, educational resources, meetings, and programs that improve public health through the practice and science of endocrinology.

Clinical Practice Chair, ENDO 2026
Ana Paula Abreu Metzger, MD, PhD

The statements and opinions expressed in this publication are those of the individual authors and do not necessarily reflect the views of the Endocrine Society. The Endocrine Society is not responsible or liable in any way for the currency of the information, for any errors, omissions or inaccuracies, or for any consequences arising therefrom. With respect to any drugs mentioned, the reader is advised to refer to the appropriate medical literature and the product information currently provided by the manufacturer to verify appropriate dosage, method and duration of administration, and other relevant information. In all instances, it is the responsibility of the treating physician or other health care professional, relying on independent experience and expertise, as well as knowledge of the patient, to determine the best treatment for the patient.

Copyright © 2026 by the Endocrine Society, 2055 L Street, NW, Suite 600, Washington, DC 20036. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, posted on the internet, or transmitted in any form, by any means, electronic, mechanical, photocopying, recording, or otherwise, without written permission of the publisher.

Requests for permissions for reproduction should be directed to Copyright Clearance Center (CCC) at www.copyright.com. For more information or to purchase copies, please contact Society Services by telephone at 202-971-3646, fax at 202-736-9704, or email at info@endocrine.org or visit the online store: www.endocrine.org/store.

ISBN: 978-1-936704-72-9
eISBN: 978-1-936704-73-6

ENDO 2026 CONTENTS



ADIPOSE TISSUE, APPETITE, OBESITY, AND LIPIDS

Lipedema at the Intersection of Endocrinology and Adipose Biology 2

José O. Alemán-Díaz, MD, PhD

GLP-1–Based Therapies: Preparing Patients With Obesity for Long-Term Maintenance and Transition 10

W. Timothy Garvey, MD

Navigating Obesity Management in Menopause and Midlife: Endocrine Mechanisms and Clinical Management 22

George Mastorakos, MD, DmedSc, MSc, Harilena Tsourouktsoglou, MD, and Georgios Valsamakis, MD, PhD

Dietary Patterns and Eating Strategies in Obesity Care: Timing, Composition, and Personalized Approaches 33

Kalpana Muthusamy, MBBS, MD

ADRENAL

New Management Options for Adults With Congenital Adrenal Hyperplasia 42

Richard J. Auchus, MD, PhD, and Benjamin Wilson, BA

Adrenal Aftermath: Postoperative Adrenal Insufficiency and Glucocorticoid Withdrawal Syndrome After Surgery for Cushing Syndrome 56

Guido Di Dalmazi, MD

Metabolic Implications of Adrenal Incidentalomas: When Do We Intervene? 63

John Newell-Price, MA, PhD

Challenges in the Diagnosis and Management of Adrenal Insufficiency 69

Richard J. M. Ross, MBBS, MD, and Miguel Debono, MD, PhD

BONE AND MINERAL METABOLISM

Beyond Basic Dual-Energy X-Ray Absorptiometry 78

Angela M. Cheung, MD, PhD, CCD

Osteoanabolic Therapies in Osteoporosis: When to Build Bone and How to Sequence Treatment 83

Christian M. Girgis, MBBS, PhD

Osteoporosis in Women With Breast Cancer	95
Peyman Hadji, MD	
PTH Replacement Therapy for Chronic Hypoparathyroidism in Adults	103
Dalal S. Ali, MD, MSc, and Aliya A. Khan, MD	
Hypophosphatasia in Adults: Recognizing, Diagnosing, and Treating a Multisystem Disease	110
Juan Carl Pallais, MD, MPH	

CARDIOVASCULAR ENDOCRINOLOGY

Interpreting Aldosterone and Renin: Tricks of the Trade	122
David Suphadetch Leungsuwan, MBBS, MMed, and Troy H. Puar, MBBS, PhD	
Beyond Statin Therapy for Hypercholesterolemia in Adults	135
Vinaya Simha, MD	
Evaluation and Management of Primary Aldosteronism	143
Anand Vaidya, MD, MMSc	

DIABETES AND VASCULAR DISEASE

Updates in the Management of Diabetic Kidney Disease	156
Rodica Busui, MD, PhD	
Type 2 Diabetes in Preconception, Pregnancy, and Postpartum	161
Aoife Maria Egan, MB BCh, BAO, PhD	
Transition of Care in Patients With Youth-Onset Type 2 Diabetes	170
Erin Finn, MD	
Use of Adjunctive Glucose-Lowering Therapies to Target Cardiometabolic Risk in Type 1 Diabetes	179
Jerry R. Greenfield, MD, PhD	
Managing Hyperglycemia in Hospitalized Patients	190
Oscar L. Morey-Vargas, MD	
GLP-1 Receptor Agonists and Pancreatitis and Pancreatic Lesions: What Is the Evidence and When Are They Contraindicated?	200
Pankaj Shah, MD	

GENERAL ENDOCRINOLOGY

Hormone Balancing: Addressing Social Media Myths	210
Michael T. McDermott, MD	

NEUROENDOCRINOLOGY AND PITUITARY

Ectopic Cushing Syndrome	222
Irina Bancos, MD, MSc	
Neuroendocrine Adverse Effects of Cancer Treatment: Immune Checkpoint Inhibitors ..	227
Alexander Faje, MD	
Current Approach to Acromegaly Care	236
Lisa Nachtigall, MD	
Management of Pituitary Apoplexy	246
Georgia Ntali, MBChB, MSc, PhD, and Niki Karavitaki, MBChB, MSc, PhD	
Diagnosis and Management of Silent Pheochromocytoma and Paraganglioma	252
Christina Pamporaki, MD, PhD	
A Structured Diagnostic Approach to Patients With Hypotonic Polyuria	262
Beatriz Santana Soares, MD, PhD, and Juliana Beaudette Drummond, MD, PhD	

PEDIATRIC AND ADOLESCENT ENDOCRINOLOGY

Management of 46,XY Difference of Sex Development Across the Lifespan: A Multidisciplinary Team Perspective	276
Berenice B. Mendonca, MD, and Sorahia Domenice, MD	
Management of Hyperphagia in Prader-Willi Syndrome and Other Monogenic Obesity Syndromes	289
Jennifer Miller, MD, MS	
Pituitary Tumor Survivorship: Transitioning From Pediatric to Adult Endocrine Care	297
Kevin C. J. Yuen, MD	

REPRODUCTIVE ENDOCRINOLOGY

Endocrine and Fertility Implications After Anabolic Steroids, Hormonal Contraception, and Gender-Affirming Hormone Therapy	310
Martin den Heijer, MD, PhD	
Update on the Management of Androgen Insensitivity Syndromes	316
Anna Nordenström, MD, PhD	

THYROID BIOLOGY AND CANCER

Recent Advances in the Management of Anaplastic Thyroid Cancer	326
Maria E. Cabanillas, MD	

Evaluation and Management of Thyroid Dysfunction in Pregnancy	336
Sun Y. Lee, MD, MSc	
Novel Insights in the Treatment of Hyperthyroidism	345
Robin Peeters, MD, PhD	
Redefining the Role of Radioactive Iodine in Differentiated Thyroid Carcinoma	353
Mabel Ryder, MD	
Targeted Therapy in Advanced Thyroid Cancer	361
Lori J. Wirth, MD	

TUMOR BIOLOGY

Management of Duodenopancreatic Neuroendocrine Tumors in Patients With Multiple Endocrine Neoplasia Type 1	372
Thorvardur (Thor) R. Halfdanarson, MD, and Omair A. Shariq, MBBS, PhD	

APPENDIX

Color Gallery	CG1
----------------------------	------------

ENDO 2026 FACULTY



2026 Endocrine Case Management: Meet the Professor Faculty

José O. Alemán-Díaz, MD, PhD
AdventHealth Orlando

Dalal S. Ali, MD, MSc
McMaster University

Richard J. Auchus, MD, PhD
University of Michigan

Irina Bancos, MD, MSc
Mayo Clinic

Rodica Busui, MD, PhD
Oregon Health and
Science University

Maria E. Cabanillas, MD
The University of Texas MD
Anderson Cancer Center

Angela M. Cheung, MD, PhD, CCD
University of Toronto

Miguel Debono, MD, PhD
University of Sheffield

Martin den Heijer, MD, PhD
Amsterdam UMC

Guido Di Dalmazi, MD
University of Bologna

Sorahia Domenice, MD
University of São Paulo

**Juliana Beaudette
Drummond, MD, PhD**
Federal University of Minas Gerais

Aoife Maria Egan, MB BCH, BAO, PhD
Mayo Clinic

Alexander T. Faje, MD
Massachusetts General Hospital

Erin Finn, MD
University of Colorado Anschutz

W. Timothy Garvey, MD
University of Alabama
at Birmingham

Christian Girgis, MBBS, PhD
Westmead Hospital and The
University of Sydney

Jerry R. Greenfield, MD, PhD
Victor Chang Cardiac Research
Institute, St. Vincent's Hospital, and
University of New South Wales

Peyman Hadji, MD
Frankfurt am Main and Philipps-
University of Marburg

**Thorvardur (Thor) R.
Halfdanarson, MD**
Mayo Clinic

Niki Karavitaki, MBChB, MSc, PhD
University of Birmingham and
Queen Elizabeth Hospital

Aliya A. Khan, MD
McMaster University

Sun Y. Lee, MD, MSc
Boston University Chobanian and
Avedisian School of Medicine

**David Suphadetch
Leungsuwan, MBBS, MMed**
Changi General Hospital

**George Mastorakos,
MD, DmedSc, MSc**
National and Kapodistrian
University of Athens

Michael T. McDermott, MD
University of Colorado
School of Medicine

Berenice B. Mendonca, MD
University of São Paulo

Jennifer L. Miller, MD, MS
University of Florida

Oscar L. Morey-Vargas, MD
Cleveland Clinic

Kalpana Muthusamy, MBBS, MD
Mayo Clinic

Lisa Nachtigall, MD
Mass General Brigham and
Harvard Medical School

John Newell-Price, MD, PhD
University of Sheffield

Anna Nordenström, MD, PhD
Karolinska Institute and
Karolinska University Hospital

Georgia Ntali, MBChB, MSc, PhD
Evangelismos Hospital,
University of Birmingham, and
Queen Elizabeth Hospital

Juan Carl Pallais, MD, MPH
Brigham and Women's Hospital,
Mass General Brigham, and
Harvard Medical School

Christina Pamporaki, MD, PhD
University Hospital Carl
Gustav Carus TU Dresden

Robin Peeters, MD, PhD
Erasmus Medical Center, Rotterdam

Troy H. Puar, MBBS, PhD
Changi General Hospital

Richard J. M. Ross, MBBS, MD
University of Sheffield

Mabel Ryder, MD
Mayo Clinic

Pankaj Shah, MD
Mayo Clinic

Omar A. Shariq, MBBS, PhD
Mayo Clinic

Vinaya Simha, MD
Mayo Clinic

Beatriz Santana Soares, MD, PhD
Federal University of Minas Gerais

Harilena Tsourouktoglou, MD
National and Kapodistrian
University of Athens

Anand Vaidya, MD, MSc
Mass General Brigham and
Harvard Medical School

Georgios Valsamakis, MD, PhD
National and Kapodistrian
University of Athens

Benjamin Wilson, BA
Boston University

Lori J. Wirth, MD
Mass General Brigham Cancer
Institute and Harvard Medical School

Kevin C. J. Yuen, MD
University of Arizona College
of Medicine and Creighton
School of Medicine

Annual Meeting Steering Committee (AMSC)

Barbara Gisella Carranza Leon, MD
AMSC Chair
Vanderbilt University Medical Center

Ana Paula Abreu Metzger, MD, PhD
Clinical Practice Chair
Brigham and Women's Hospital

Raghavendra Mirmira, PhD, MD
Clinical Science Chair
University of Chicago

Erik Nelson, BSc, PhD
Basic Science Chair
University of Illinois at
Urbana-Champaign

Annual Meeting Steering Committee Clinical Peer Reviewers

Olga Astapova, MD, PhD

Laura Boucai, MD

Ana Pinheiro Machado
Canton, MD, PhD

Diane Donegan, MBBCh BAO, MRCPI

Christa E. Flueck, MD

Oksana Hamidi, DO

Edward Chiaming Hsiao, MD, PhD

Hebatullah M. Ismail, MD

Suzanne Jan De Beur, MD

Peter Kamenicky, MD, PhD

Srividya Kidambi, MD

Katja Kiseljak-Vassiliades, DO

Connie Baum Newman, MD

Theodora Pappa, MD, PhD

Elizabeth N. Pearce, MD, MSc

Luca Persani, MD, PhD

Julie Refardt, MD, PhD

Laura Torchen, MD

Elena Tsourdi, MD

Adina Turcu, MD, MS

Jun Yang, MBBS, PhD

Corrine Welt, MD

ENDO 2026 OVERVIEW



OVERVIEW

Endocrine Case Management: Meet the Professor is designed to provide physicians with a concise, high-quality review of 40 common and rare endocrine disorders to help you keep your practice current. It features case-based clinical vignettes and rationales written by experts across all areas of endocrinology, diabetes, and metabolism.

ACCREDITATION STATEMENT

The Endocrine Society is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education (CME) for physicians. The Endocrine Society has received Accreditation with Commendation.



The Endocrine Society designates this enduring material for a maximum of 30 *AMA PRA Category 1 Credits*[™]. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

MAINTENANCE OF CERTIFICATION

Successful completion of this CME activity, including participation in the evaluation component, enables the participant to earn up to 30 MOC points in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. It is the CME activity provider's responsibility to submit participant completion information to the Accreditation Council for Continuing Medical Education for the purpose of granting ABIM MOC credit.



CME credits and/or MOC points for the activities related to this material must be claimed by December 31, 2028. For questions about content or obtaining CME credit or MOC points, please contact the Endocrine Society at info@endocrine.org.

LEARNING OBJECTIVES

Endocrine Case Management: Meet the Professor enables learners to assess their knowledge of all aspects of endocrinology, diabetes, and metabolism.

Upon completion of this educational activity, learners will be able to:

- Recognize the clinical manifestations of endocrine and metabolic disorders and select among current diagnostic, management, and therapeutic options.
- Identify risk factors for endocrine and metabolic disorders and develop prevention strategies.
- Evaluate endocrine and metabolic manifestations of systemic disorders.
- Use existing resources on clinical guidelines and treatment recommendations for endocrine and related metabolic disorders to guide diagnosis and treatment.

TARGET AUDIENCE

Endocrine Case Management: Meet the Professor provides case-based education for clinicians seeking to improve patient care.

STATEMENT OF INDEPENDENCE

As a CME provider accredited by the Accreditation Council for Continuing Medical Education, the Endocrine Society has a policy to ensure that the content of this educational activity is high-quality, balanced, independent, objective, and scientifically rigorous. The scientific content of this activity was developed under the supervision of the Endocrine Society's Annual Meeting Steering Committee.

DISCLOSURE POLICY

The faculty, committee members, and staff who are in a position to control the content of this activity are required to disclose to the Endocrine Society and to learners any relevant financial relationship(s) of the individual or spouse/partner that have occurred within the last 12 months with any commercial interest(s) whose products or services are related to the content. Financial relationships are defined as remuneration in any amount from the commercial interest(s) in the form of grants; research support; consulting fees; salary; ownership interest (e.g., stocks, stock options, or ownership interest excluding diversified mutual funds); honoraria or other

payments for participation in speakers' bureaus, advisory boards, or boards of directors; or other financial benefits. The intent of this disclosure is not to prevent planners with relevant financial relationships from planning or delivering content, but rather to provide learners with information that enables them to make their own judgments about whether these financial relationships may have influenced the educational activity's exposition or conclusions. The Endocrine Society has reviewed all disclosures and resolved or managed all identified conflicts of interest, as applicable.

The Endocrine Society has reviewed these relationships to determine which are relevant to this activity's content and has resolved any identified conflicts of interest for these individuals.

DISCLOSURE POLICY

The faculty reported the following relevant financial relationship(s) during the content development process for this activity:

Richard J. Auchus, MD, PhD Adrenas (Contracted Research), Aspect Biosystems (Consultant), Crinetics (Consultant and Contracted Research), Lundbeck (Consultant), Lundbeck (Contracted Research), Merck (Consultant), Neurocrine (Consultant and Contracted Research), OMass (Consultant), Vertex (Consultant).

Rodica Busui, MD, PhD Novo Nordisk (Clinical Trial Investigator and Consultant), Averitas Pharma (Consultant), Bayer Inc. (Clinical Trial Investigator and Consultant), Lexicon (Clinical Trial Investigator and Consultant), Roche Diagnostics (Consultant), and Vertex (Consultant).

Maria E. Cabanillas, MD Bayer Inc. (Consultant), EISAI (Research funding), Exelixis (Consultant and Research Funding), Genentech (Research Funding), Lilly (Consultant), Merck (Research Funding), and Novartis (Consultant).

W. Timothy Garvey, MD Novo Nordisk (Advisory Board and Clinical Trial Investigator), AbbVie (Advisory Board), Boehringer-Ingelheim (Advisory Board), Corcept Therapeutics (Advisory Board), Eli Lilly & Company (Advisory Board and Clinical Trial Investigator), Fractyl Health (Advisory Board), Gan & Lee (Advisory Board), Graviton Biosciences (Advisory Board), i2o Therapeutics (Advisory Board), Kaleira (Clinical Trial Investigator), Keros Therapeutics (Advisory Board), Madrigal (Advisory Board), Neurocrine Biosciences (Advisory Board), Pfizer (Advisory Board), Regeneron (Advisory Board), Roche Pharmaceuticals (Advisory Board and Clinical Trial Investigator), TERNs Pharmaceuticals (Advisory Board and Clinical Trial Investigator), Viking Therapeutics (Clinical Trial Investigator), and Zealand (Advisory Board and Clinical Trial Investigator).

Christian Girgis, MBBS, PhD Theramex (Advisory Board).

Jerry R. Greenfield, MBBS, PhD Novo Nordisk (Recipient of drug and placebo for a study of semaglutide in type 1 diabetes).

Thorvardur R. Halfdanarson, MD Biomea Fusion (Advisory Board), Boehringer Ingelheim (Advisory Board), Camurus (Clinical Trial Investigator), Crinetics (Clinical Trial Investigator), Curium (Advisory Board), Exelixis (Advisory Board and Clinical Trial Investigator), ITM (Clinical Trial Investigator), Novartis (Clinical Trial Investigator), Perspective Therapeutics (Clinical Trial Investigator), RayzeBio (Clinical Trial Investigator), Rezolute (Clinical Trial Investigator), Sanofi (Advisory Board), and Thermo Fisher (Clinical Trial Investigator).

Aliya A. Khan, MD Ascendis Pharma (Grant Recipient).

Sun Y. Lee, MD, MSc Alcimed (Consultant), American Association of Clinical Endocrinology (Honorarium Recipient), Florida Endocrinology Association (Honorarium Recipient), and Guidepoint (Consultant).

Jennifer L. Miller, MD, MS Rhythm Therapeutics (Clinical Trial Investigator) and Soleno Therapeutics (Clinical Trial Investigator).

Oscar L. Morey-Vargas, MD Asofarma (Teaching and Speaking).

Lisa Nachtigall, MD Camurus (Advisory Board), Chiesi (Advisory Board, Consultant, and Contracted Research), Crinetics (Advisory Board and Consultant), and Recordati Rare Disease (Advisory Board and Grant Recipient).

John Newell-Price, MD, PhD Crinetics (Advisory Board), Recordati (Writing Committee), and Sparrow Therapeutics (Grant Recipient).

Juan Carl Pallais, MD, MPH Alesta Therapeutics (Consultant), Alexion Pharmaceuticals Inc. (Grant Recipient), Alexion Pharmaceuticals Inc. (Honorarium Recipient), Ensho Health (Advisory Board), Soft Bones (Speakers' Bureau), UpToDate (Author).

Troy H. Puar, MBBS, PhD AstraZeneca (Clinical Trial and Travel Grant).

Richard J. M. Ross, MBBS, MD Neurocrine (Consultant).

Anand Vaidya, MD, MMSc AstraZeneca (Consultant), Corcept (Consultant), Doris Duke Charitable Foundation (Grant Recipient), Mineralys (Advisory Board), and NIH (Grant Recipient).

Benjamin Wilson, BA Neurocrine Biosciences (Consultant).

Lori J. Wirth, MD AbbVie Inc (Advisory Board), Bicara (Advisory Board), Coherus (Advisory Board), EISAI (Consultant), Fore Biopharma (Advisory Board), Leo Pharma (Advisory Board), Novartis (Consultant), Pyxis (Advisory Board), and Tubulis (Advisory Board).

Kevin C. J. Yuen, MD Ascendis Pharma (*Advisory Board and Contracted Research*), Camurus (*Advisory Board*), Chiesi (*Advisory Board*), Crinetics (*Advisory Board*), Neurocrine (*Advisory Board and Speakers' Bureau*), Novo Nordisk (*Advisory Board, Contracted Research, and Speakers' Bureau*), Recordati (*Advisory Board and Speakers' Bureau*), and Sparrow Therapeutics (*Contracted Research*).

The following faculty reported no relevant financial relationships:

José O. Alemán-Díaz, MD, PhD; Dalal S. Ali, MD, MSc; Irina Bancos, MD, MSc; Angela M. Cheung, MD, PhD, CCD; Miguel Debono, MD, PhD; Martin den Heijer, MD, PhD; Guido Di Dalmazi, MD; Sorahia Domenice, MD; Juliana Beaudette Drummond, MD, PhD; Aoife Maria Egan, MB BCh, BAO, PhD; Alexander T. Faje, MD; Erin Finn, MD; Peyman Hadji, MD; Niki Karavitaki, MBChB, MSc, PhD; David Suphadetch Leungsuwan, MBBS, MMed; George Mastorakos, MD, DmedSc, MSc; Michael T. McDermott, MD; Berenice B. Mendonca, MD; Kalpana Muthusamy, MBBS, MD; Anna Nordenström, MD, PhD; Georgia Ntali, MBChB, MSc, PhD; Christina Pamporaki, MD, PhD; Robin Peeters, MD, PhD; Mabel Ryder, MD; Pankaj Shah, MD; Omair A. Shariq, MBBS, PhD; Vinaya Simha, MD; Beatriz Sanatana Soares, MD, PhD; Harilena Tsourouktoglou, MD; and Georgios Valsamakis, MD, PhD.

The following AMSC peer reviewers reported relevant financial relationships:

Barbara Gisella Carranza Leon, MD Novartis, IONIS Pharmaceutical Inc, NIH, FH Foundation, Regenzbio, Inc (*Coinvestigator*), Maintenance of Certification Committee: American Board of Obesity Medicine (*Member*).

Diane Donegan, MB BCh BAO, MRCPI Corcept, Recordati (*Investigator*); Camarus (*Advisory Board*).

Christa E. Flueck, MD European Society Paediatric Endocrinology (*Advisory Board*); Novo Nordisk, Merck, Pfizer, Sandoz, Swiss National Science Foundation, European Society Paediatric Endocrinology (*Grant Recipient*); Novo Nordisk, Pfizer, EffRx (*Consultant*); Sandoz (*Speaker*); *Hormone Research in Paediatrics* (*Editor*).

Oksana Hamidi, DO Corcept Therapeutics, Crinetics Pharmaceuticals, Neurocrine Biosciences, Recordati Rare Diseases, Camurus (*Advisory Board*); Xeris Pharma (*Speaker*).

Edward Chiaming Hsiao, MD, PhD International FOP Association, FD/MAS Alliance, International Clinical Council on FOP (*Advisory Board*); *Endocrine Journal*, *Journal of Bone and Mineral Research* (*Editor*); Clementia Pharmaceuticals/Ipsen Pharmaceuticals, Ultragenyx (*Research Investigator*).

Suzanne Jan De Beur, MD American Society of Bone and Mineral Research (*Past President*); XLH Network, Current Osteoporosis Reports, Ultragenyx, Kyowa Kirin, Ascendis (*Advisory Board*); *Bone Reports* (*Editorial Board*); Mereo, Kyowa Kirin, Calcilytics (*Researcher*); NIH, FDA (*Grant Reviewer*); PCORI (*Investigator*).

Peter Kamenicky, MD, PhD Inserm (*Member of Commission Scientifique Spécialisée CSS3*); Pfizer (*Speaker*); Amolyt Pharma (*Investigator and Speaker*); Ascendis Pharma (*Advisory Board*); and Calcilytix (*Investigator*).

Srividya Kidambi, MD TOPS Inc (*Medical Editor and Director*).

Katja Kiseljak-Vassiliades, DO HRA Pharma (*Advisory Board*).

Raghavendra Mirmira, PhD, MD *Journal of Clinical Endocrinology and Metabolism* (*Editor*); Veralox Therapeutics (*Advisory Board*); Veralox Therapeutics, HiberCell, Inc (*Investigator-Initiated Award*).

Connie Baum Newman, MD Food and Drug Association (FDA) for Endocrinologic and Metabolic Diseases (*Advisory Committee and Consultant*); Federation of Medical Women of Canada (*Honorarium*).

Theodora Pappa, MD, PhD Endocrine Fellows Foundation (*Board of Directors*), *Frontiers* (*Editorial Board Member*), *Clinical Thyroidology* (*Associate Editor*), and *Eisai* (*Consultant*).

Elizabeth N. Pearce, MD, MSc Iodine Global Network (*Management Council*); *Journal of Clinical Endocrinology and Metabolism*, DynaMed-thyroid (*Editor*); National Dairy Council, Merck China Symposium (*Speaker*); American Thyroid Association (*Guidelines Task Force*); World Health Organization (*Advisory Panel*).

Luca Persani, MD, PhD Merck, Egetis, Sandoz (*Advisory Board*); Merck (*Speaker*); *Journal of Clinical Endocrinology and Metabolism*, European Thyroid Association, *European Thyroid Journal*, Pituitary Society, *Pituitary* (*Editor*).

Laura Torchen, MD Levo therapeutics, IBSA Pharma (*Investigator*).

Elena Tsourdi, MD Amgen, Kyowa Kirin, UCB, Alexion, Ascendis (*Clinical Studies*); Amgen, UCB, Ascendis (*Advisory Boards and Lectures*); UCB, Takeda, Ascendis (*Educational Grants*).

Adina Turcu, MD, MS NIH/NIDDK/NHLBI (*Grant Recipient*), ESAP (*Questions Writer*), *JCEM: Case Reports* (*Deputy Editor*), and AstraZeneca (*Advisor*).

Corrine Kolka Welt, MD UpToDate.

The following AMSC members reported no relevant financial relationships:

Ana Paula Abreu Metzger, MD, PhD; Olga Astapova, MD, PhD; Laura Boucai, MD; Ana Pinheiro Machado Canton, MD, PhD; Hebatullah M. Ismail, MD; Julie Refardt, MD, PhD; and Jun Yang, MBBS, PhD.

DISCLAIMERS

The information presented in this activity reflects the faculty's opinions and is not necessarily the official position of the Endocrine Society.

USE OF PROFESSIONAL JUDGMENT:

The educational content in this activity covers basic principles of diagnosis and therapy and does not replace individual patient assessment based on the health care provider's examination of the patient and on consideration of laboratory data and other patient-specific factors. Medical standards change as new data become available.

DRUGS AND DOSAGES:

When prescribing medications, the physician is advised to review the product information sheet for each drug to verify the conditions of use and to identify any changes to the drug's dosage schedule or contraindications.

POLICY ON UNLABELED/OFF-LABEL USE

The Endocrine Society has determined that disclosure of unlabeled, off-label, or investigational use of commercial products is informative for audiences and therefore requires that this information be disclosed to learners at the beginning of the presentation. Uses of specific therapeutic agents, devices, and other products discussed in this educational activity may not be the same as those indicated in product labeling approved by the Food and Drug Administration (FDA). The Endocrine Society requires that any discussions of such "off-label" use be based on scientific research that conforms to generally accepted standards of experimental design, data collection, and data analysis. Before recommending or prescribing any therapeutic agent or device, learners should review the complete prescribing information, including indications, contraindications, warnings, precautions, and adverse events.

PRIVACY AND CONFIDENTIALITY STATEMENT

The Endocrine Society will record learners' personal information, as provided in CME evaluations, to issue and track CME certificates. The Endocrine Society may also track aggregate responses to questions in activities and evaluations and use these data to inform the ongoing evaluation and improvement of its CME program. No individual performance data or other personal information collected from evaluations will be shared with third parties.

ACKNOWLEDGMENT OF COMMERCIAL SUPPORT

The activity is not supported by educational grants or other funds from any commercial supporters.

AMA PRA CATEGORY 1 CREDIT™ (CME) INFORMATION

To receive a maximum of 30 *AMA PRA Category 1 Credits™*, participants must complete the online interactive module and activity evaluation at the Endocrine Society's Center for Learning: <https://education.endocrine.org>. Participants must achieve a minimum score of 70% to claim CME credit. After completing the module, if participants do not achieve a minimum score of 70%, they may revise their answers and make additional attempts to reach a passing score. Learners also have the option to clear all answers and start over.

METHOD OF PARTICIPATION

This enduring material is presented online and in print format. The estimated time to complete this activity, including review of material, is 30 hours. Participants must achieve a minimum score of 70% to claim CME credit and MOC points. After completing the module, if participants do not achieve a minimum score of 70%, they may revise their answers and make additional attempts to reach a passing score. Participants also have the option to clear all answers and start over.

SYSTEM REQUIREMENTS

To complete this activity, participants must have access to a computer or mobile device with an Internet connection and use an up-to-date version of a major Web browser, such as Internet Explorer 10+, Firefox 32+, Safari, or Google Chrome 37+. In addition, cookies and JavaScript must be enabled in the browser's settings.

LAST REVIEW: May 2026

ACTIVITY RELEASE: June 13, 2026

ACTIVITY EXPIRATION DATE: December 31, 2028 (date after which this enduring material is no longer certified for *AMA PRA Category 1 Credits™* and ABIM Medical Knowledge MOC points)

For questions about content or obtaining CME credit or MOC points, please contact the Endocrine Society at info@endocrine.org.

COMMON ABBREVIATIONS IN MEET THE PROFESSOR

ACTH = corticotropin	FSH = follicle-stimulating hormone	NPH insulin = neutral protamine Hagedorn insulin
ACE inhibitor = angiotensin-converting enzyme inhibitor	GH = growth hormone	PCSK9 inhibitor = proprotein convertase subtilisin/kexin 9 inhibitor
ALT = alanine aminotransferase	GHRH = growth hormone-releasing hormone	PET = positron emission tomography
AST = aspartate aminotransferase	GLP-1 receptor agonist = glucagonlike peptide 1 receptor agonist	PSA = prostate-specific antigen
BMI = body mass index	GnRH = gonadotropin-releasing hormone	PTH = parathyroid hormone
CNS = central nervous system	hCG = human chorionic gonadotropin	PTHrP = parathyroid hormone-related protein
CT = computed tomography	HDL = high-density lipoprotein	SGLT-2 inhibitor = sodium-glucose cotransporter 2 inhibitor
DHEA = dehydroepiandrosterone	HIV = human immunodeficiency virus	SHBG = sex hormone-binding globulin
DHEA-S = dehydroepiandrosterone sulfate	HMG-CoA reductase inhibitor = 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor	T₃ = triiodothyronine
DNA = deoxyribonucleic acid	IGF-1 = insulinlike growth factor 1	T₄ = thyroxine
DPP-4 inhibitor = dipeptidyl-peptidase 4 inhibitor	LDL = low-density lipoprotein	TPO antibodies = thyroperoxidase antibodies
DXA = dual-energy x-ray absorptiometry	LH = luteinizing hormone	TRH = thyrotropin-releasing hormone
FDA = Food and Drug Administration	MCV = mean corpuscular volume	TRAb = TSH-receptor antibodies
FGF-23 = fibroblast growth factor 23	MIBG = <i>meta</i> -iodobenzylguanidine	TSH = thyrotropin
FNA = fine-needle aspiration	MRI = magnetic resonance imaging	VLDL = very low-density lipoprotein



ADIPOSE TISSUE, APPETITE, OBESITY, AND LIPIDS



ADIPOSE TISSUE, APPETITE, OBESITY, AND LIPIDS

Lipedema at the Intersection of Endocrinology and Adipose Biology

José O. Alemán-Díaz, MD, PhD

Figure. Stages of Lipedema According to Progression of Fat Accumulation and Changes to Skin and Lymphatic System 3



Adapted from the Lipedema Foundation, lipedema.org

Lipedema at the Intersection of Endocrinology and Adipose Biology

José O. Alemán-Díaz, MD, PhD. Laboratory of Translational Obesity Research, Translational Research Institute, AdventHealth Orlando, Orlando, Florida; Email: jose.alemandiaz@adventhealth.com

Educational Objectives

After reviewing this chapter, learners should be able to:

- Recognize lipedema as a distinct disorder of loose connective tissue under the umbrella of obesity, distinct from vascular disease–related lymphedema.
 - Describe endocrinology and adipose biology relevant to the development and future treatment of lipedema.
 - Discuss current and emerging treatment options to reduce morbidity from lipedema.
-

Significance of the Clinical Problem

What Is Lipedema?

Lipedema is a chronic disorder of subcutaneous adipose tissue characterized by disproportionate fat accumulation in the limbs (usually the legs, sometimes the arms) and almost exclusively affects women. First described in 1940 by Allen and Hines, lipedema is distinct from general obesity and lymphedema, with morbidity decreasing quality of life.^{1,2} In lipedema, excessive fat deposition causes symmetric enlargement of the legs and/or arms, often presenting as a columnlike leg shape, while hands and feet remain unaffected until late stages. A hallmark is that the affected fat

tissue is painful to palpation and prone to bruising with minor trauma, indicative of capillary fragility. Unlike common obesity, the fat in lipedema is diet-resistant, meaning patients often report little improvement in limb size even after significant weight loss. This can help distinguish lipedema from simple obesity. Importantly, lipedema fat deposition is usually bilateral and symmetric, differentiating it from lymphedema, in which swelling is often unilateral.³ In summary, lipedema is recognized as a loose connective tissue disease of fat, involving nodular, fibrotic fat accumulation on the hips, buttocks, and limbs, arising in the context of hormonal and developmental triggers (eg, puberty), and causing progressive disproportionate limb enlargement with pain.

Diagnostic Criteria

The diagnosis of lipedema is clinical, based on a thorough patient history, characteristic physical exam findings, and exclusion of other conditions. Formal diagnostic criteria were established by the US Standard of Care consensus,⁴ which outlined key features for identifying lipedema in practice. There is no specific laboratory test or biomarker for lipedema; instead, clinicians rely on recognizing a constellation of signs and symptoms. Major criteria include: (1) a bilateral, symmetric distribution of abnormal subcutaneous fat, typically affecting the legs (from the hips down to the ankles) and sometimes the arms, with sparing of the feet and hands (producing an

abrupt cutoff or “cuffing” at the ankles or wrists); (2) a negative Stemmer sign, meaning the skin at the base of the toes and fingers can be pinched and lifted (in contrast to lymphedema, where this is usually impossible due to swelling); (3) little or no pitting edema on pressure, at least in early stages (any swelling in lipedema is nonpitting and generally minimal unless lymphatic compromise develops). Additional supportive criteria include pain, tenderness, and easy bruising in the affected fat tissue; patients often report that even slight pressure or minor bumps cause pain and bruising in the legs. Onset or worsening during hormonal changes, such as around puberty or pregnancy, is another common historical clue. Importantly, disproportionate fat distribution is a key feature; the lower body is much more enlarged than the upper body, sometimes accompanied by palpable fat nodules in the subcutis. In contrast, the trunk remains comparatively slender, and the feet are not swollen, distinguishing lipedema from generalized obesity or lymphedema. Patients typically do not experience significant

improvement in limb size with calorie restriction or exercise (diet-resistant fat), which can aid in diagnosis. Clinical examination may also reveal textural changes in the skin, depending on the stage of the disease. In early stages, the skin may feel rubbery with fine nodularity, progressing to larger nodules and an irregular “mattress-like” texture in later stages as fibrotic nodules form. Formal staging systems grade the skin and tissue nodularity: stage 1, smooth skin, soft nodules; stage 2, uneven skin surface, large nodules (apple- to walnut-sized), fat thickening, and fibrosis; stage 3, large extrusion of fat tissue, indurated masses and lobules of fat; and stage 4, sometimes used to denote lipedema with accompanying lymphedema (lipo-lymphedema), as shown in the *Figure*. Additionally, lipedema is classified by type based on regional distribution: type I: buttocks/hips; type II: buttocks to the knees; type III: down to the ankles; type IV: arms; type V: calves. These staging classifications, illustrated in the *Figure*, help document the extent of disease for clinical and research purposes.

Figure. Stages of Lipedema According to Progression of Fat Accumulation and Changes to Skin and Lymphatic System



Adapted from the Lipedema Foundation, lipedema.org

[Color—Print (Color Gallery page CG3) or web & ePub editions]

Prevalence

Lipedema is believed to be far more common than historically recognized, affecting a substantial percentage of the female population. Precise prevalence figures remain uncertain because of underdiagnosis and past confusion with obesity, but recent estimates suggest that roughly 11% (up to nearly 1 in 9) of adult women worldwide may have lipedema. In specific settings, even higher rates have been reported; for example, a German literature review reported that 6% to 8% of women in the general population could have lipedema, and 15% to 19% of women seen in specialty vascular clinics were diagnosed with lipedema.⁵ In contrast, lipedema is exceedingly rare in men, with only isolated case reports, usually in men with extreme hormonal disorders, such as testosterone deficiency or estrogen excess, or liver disease.

Practice Gaps

- Lipedema is common among women seeking care for weight management or vascular medicine.
- Lipedema does not respond to medical or surgical weight-loss interventions. Cornerstones of therapy include compression and, in advanced cases, plastic surgery to remove affected adipose tissue.
- As a disorder of loose connective tissue, including adipose tissue, lipedema has important hormonal connections to its development and pathophysiology that endocrinologists should be aware of.

Discussion

Pathophysiology

The pathophysiology of lipedema is complex and generally involves abnormal expansion of subcutaneous adipose tissue, associated microvascular changes influenced by genetic predisposition, and environmental cues. Conceptually, this pathophysiology can be

organized into vascular, endocrine, and adipose components, among others, for the purposes of this chapter.

Vascular Component

Despite the “edema” suffix, lipedema does not involve fluid accumulation in early stages. Unlike lymphedema, the hallmark of lipedema is disproportionate, symmetric accumulation of painful adipose tissue in the upper or lower extremities. Feet, hands, trunk, neck, and head are not affected.⁴ The Stemmer sign, in which the examiner cannot pinch the skin of the toes, is negative in lipedema. By contrast, lymphedema stems from impaired lymphatic capillary drainage, often begins distally in the feet, and tends to be asymmetric. In this context, lipedema is often considered to have relative lymphatic sparing and minimal edema in early stages. The microvasculature in lipedema is compromised, leading to easy bruising and increased vascular permeability in fat, both of which indicate a microangiopathic state. Leaky capillaries may lead to bruising after minor trauma, as well as to pressure, heaviness, and pain on palpation. In later stages, secondary lymphedema can coexist, resulting in the condition termed lipo-lymphedema. The International Lipedema Consensus of 2020 proposed that chronic inflammation and tissue hypoxia in the enlarging fat are likely causes of the pain in lipedema, rather than fluid edema per se.⁴ This insight shifts the treatment focus in lipedema beyond purely decompressive therapies toward anti-inflammatory and antiobesity strategies, among others. Vascular and lymphatic factors in lipedema appear secondary to adipose pathology but can progress as the disease advances, underscoring the importance of vascular and lymphatic care in later disease stages.

Endocrine Component

Lipedema development is closely associated with hormonal changes across the female life course.⁶ Lipedema often appears or worsens during periods of major hormonal flux and weight change, classically

at puberty, sometimes during pregnancy, and frequently at menopause. A common hypothesis is that lipedema may be linked to estrogen surges during these transitions. Research suggests that lipedema adipose tissue may have an abnormal estrogen profile, including an imbalance in estrogen receptor alpha ($ER\alpha$) and beta ($ER\beta$) levels. Estrogen promotes the accumulation of adipose tissue in the gluteofemoral distribution, consistent with the lipedema fat distribution. One study reported reduced $ER\alpha$ and increased $ER\beta$ in adipose tissue from patients with lipedema, which could promote lipid accumulation in lower-body depots.⁷ Progesterone and androgen levels have also been implicated in pregnancy and the menopausal transition, respectively, but direct evidence is lacking. Hormonal triggers are central to when lipedema manifests and flares, making it a quintessential example of an endocrine-influenced adipose disorder. Future research focused on the molecular endocrinology of lipedema could reveal targeted ways to interrupt this hormonal trigger cycle.

Adipose Component

At its core, lipedema is a disorder of adipose tissue within loose connective tissue. Adipocytes play the starring role in pathophysiology, with both hypertrophic and hyperplastic changes documented.^{7,8} Histological studies of lipedema fat have demonstrated adipocyte hypertrophy (enlarged fat cells) and adipocyte hyperplasia (increased number of fat cells) in the affected areas.⁹ These observations imply that the subcutaneous tissue in lipedema can expand markedly in volume, partly by each fat cell storing more lipid and swelling, and partly by recruiting or differentiating new fat cells. This contrasts with typical obesity, where hypertrophy tends to predominate.¹⁰ Lipedema fat often forms a nodular texture; microscopically, nodules of fibrotic connective tissue surround groups of adipocytes, especially in later stages. Notably, in early-stage lipedema, there may be relatively little fibrosis; the skin remains soft, and the fat, although nodular, is not hardened. Consequently, stage 1 lipedema skin feels smooth even though underlying fat nodules

can be palpated like small peas. As the condition progresses to stages 2 and 3, fibrosis increases in the subcutis: collagen deposition and scarlike tissue form around larger clumps of fat, causing the overlying skin to indent and dimple (mimicking cellulite, but more extreme). By stage 3, sizable lobules of fat develop, suspended by fibrous septa, leading to deforming overhangs of tissue around the knees or arms. Despite fibrosis of the fat in lipedema, skin elasticity often remains relatively good (until very advanced disease), which is why compression and surgery can achieve good skin retraction. Importantly, pure lipedema (stages 1-3) does not cause the dermal fibrosis or peau d'orange texture seen in lymphedema. The skin of lipedema, aside from the nodules underneath, can be pinchable and is not woody or indurated. Only when lymphedema overlaps does the skin begin to thicken. Another component of the adipose tissue change is the inflammatory cell infiltrate. Research indicates that lipedema fat depots have a distinct immune cell composition compared to both normal and obesity-related fat. There is evidence of a higher proportion of M2 macrophages (alternatively activated, anti-inflammatory macrophages) in lipedema tissue, whereas obesity-related fat tends to have more proinflammatory M1 macrophages. This M2 predominance might sound counterintuitive, since lipedema is painful (one would expect inflammation to cause pain), but it likely reflects a chronic, unresolved state of tissue remodeling. Some theorize that the pain in lipedema is due to nerve irritation within the expanding fat. Indeed, hypoxia and mild inflammation can sensitize nerves. One hypothesis is that as fat hypertrophies, small nerve fibers in the tissue are stretched or compressed, leading to a neuropathic pain element. There is also evidence of abnormal adipose stem-cell function in lipedema. A study found that adipose-derived stem cells from patients with lipedema have impaired adipogenesis *in vitro* and altered adipokine expression. Intriguingly, these differences potentially could serve as biomarkers of lipedema or even as therapeutic targets. From a life-course perspective, once lipedema fat forms,

it tends to be resistant to loss. Calorie restriction or bariatric surgery may reduce overall weight, but often spares the lipedema fat. This suggests a fundamental dysregulation of adipose metabolism in lipedema tissue. Several groups have postulated defects in catecholamine or hormone receptors on lipedema fat cells, or localized adipose hypoxia, leading to insulin resistance in that tissue, but these hypotheses remain under investigation. Finally, unlike fibrotic lymphedema tissue, lipedema adipose tissue remains “soft” fat (especially in the early stages), composed of a loose connective tissue matrix that holds water and fat; thus, the term “loose connective tissue disease” is used for lipedema. The fat is not scarred in the way of postradiation lymphedema tissue; rather, it is an overgrowth of otherwise normal fat cells and supporting stroma. Understanding the adipose component is key to developing treatments; this is why liposuction can dramatically help by physically removing pathological fat cells. In summary, lipedema involves *both* adipocyte hypertrophy and hyperplasia, leading to a significant expansion of subcutaneous fat mass with distinctive nodular fibrosis and inflammatory changes. This adipose pathology, influenced by hormones and genetics, is the central driver of the disease, around which vascular and lymphatic issues revolve.

Treatment Options

There is currently no cure for lipedema, but a variety of treatments can manage symptoms, improve function, and slow disease progression. Optimal care is usually multidisciplinary, combining medical management, nutritional guidance, physical therapies, and sometimes surgery. The overall goals are to reduce pain and swelling, restore mobility, prevent or treat lymphedema, and address the psychosocial impact. As a provider, it is important to set expectations: while treatments can significantly improve quality of life, they typically do not eliminate all lipedema tissue (except through surgical removal), and ongoing maintenance is needed. The main

treatment modalities in the standard of care are discussed in the following sections.

Medical Management

Medical management of lipedema focuses on treating symptoms and related conditions, since no pill or injection can yet specifically remove lipedema fat. Pain management is one aspect. Many patients benefit from analgesics, including nonsteroidal anti-inflammatory agents. If neuropathic pain is suspected, medications such as gabapentin or duloxetine may be tried under physician guidance. Treating coexisting issues is also crucial. For example, if venous insufficiency or varicose veins contribute to edema, medical or interventional treatments for those conditions (such as compression or vein ablation) should be provided. Likewise, managing hypothyroidism, vitamin D deficiency, or any metabolic disorders can help overall tissue health. A cornerstone of medical management is compression therapy, which is considered “medical” because it is prescribed and fitted by health care providers. Consistent use of graduated compression stockings or tights (usually class II, 20–30 mm Hg or higher) helps reduce daily swelling, supports the tissues, and can diminish pain and bruising by limiting microvascular leak. Finding the right compression garment often involves working with a specialist.¹¹ When lipedema overlaps with fluid edema, involvement of a certified lymphedema therapist is extremely beneficial. Certified lymphedema therapists are trained in complete decongestive therapy, which includes manual lymphatic drainage massage, multilayer compression bandaging, exercises, and skin care education.

Surgical Treatment

Another important treatment decision is whether to pursue surgical treatment, specifically, liposuction.⁵ Liposuction for lipedema is generally categorized as a surgical intervention, but because it is one of the few ways to actually remove the pathological fat, it is part of the standard care discussion. Typically, conservative measures are tried for at least 6 to 12 months, and if the patient

still has significant pain or functional limitation, liposuction is considered. Liposuction (ideally using the tumescent technique with microcannulas by surgeons experienced in lipedema) can dramatically improve symptoms: it permanently removes lipedema fat cells, resulting in reduced limb size, improved contour, and often much less pain and bruising. However, liposuction usually requires multiple sessions, as each limb may require 2 to 3 operations staged months apart, and it carries risks typical of any surgery, including fluid shifts and potential lymphatic injury. In experienced hands, liposuction is safe and effective. For many patients with moderate-to-severe lipedema, liposuction is the closest thing to a “definitive” treatment. Some countries, such as Germany, have officially recognized liposuction as part of lipedema therapy and even cover it under insurance in certain cases. In the United States, insurance coverage varies and can be challenging.

Nutrition

There is no single scientifically proven “lipedema diet” at this time. However, several dietary approaches have been advocated by experts and patients, including anti-inflammatory diets, lower-carbohydrate or ketogenic diets, intermittent fasting, gluten-free or allergy-elimination diets, and general weight-loss diets. The common theme across these nutritional strategies is promoting a healthy, balanced diet that the patient can sustain over the long term. In conclusion, nutrition is a cornerstone of lipedema management, aimed at optimizing body composition and reducing inflammation. As Tomada noted,⁶ proper nutrition and weight management can preserve mobility and function and potentially slow disease progression.

After decades of relative neglect, patient advocacy, scientific curiosity, and clinical urgency are driving progress in treating lipedema.¹² Ultimately, the vision for the future is one where lipedema is no longer an “orphan” condition, but a well-characterized medical entity with a range of effective treatments.

Clinical Case Vignettes

Case 1

A 52-year-old woman with class 2 obesity (BMI = 36 kg/m²) presents with worsening bilateral leg swelling. She reports menarche at age 11 and a distant history of leg swelling during puberty. She had difficulty maintaining her weight throughout young adulthood, with “big legs.” Over the past 3 years, the swelling has markedly increased, causing discomfort when putting on clothes and walking. She endorses occasional hot flashes and misses periods every 2 to 3 months.

On physical examination, the patient appears well and has a gynoid pattern of obesity. Palpation of her lower extremities reveals pearl-like nodules along her thighs and hypersensitivity to touch, with associated discomfort.

Laboratory test results are normal, including those from a basic metabolic panel, a complete blood cell count, B-type natriuretic peptide, and measurement of TSH and vitamin B₁₂.

What additional testing (if any) is needed to diagnose lipedema in this patient?

- A. No additional testing needed
- B. Vascular scintigraphy to exclude lymphedema
- C. Coagulation tests
- D. LH, FSH, and estradiol measurement

Answer: A) No additional testing needed

The diagnosis of lipedema relies on a thorough patient history, characteristic physical findings, and the exclusion of other conditions. At this time, there are no specific laboratory findings with diagnostic utility in lipedema, although exciting research is ongoing. Consequently, the lab-based findings listed in Answers B, C, and D can be suggestive but not diagnostic. Vascular scintigraphy is an invasive test that uses nuclear dyes to assess reverse flow in the lymphatic vasculature and can help exclude lymphedema. However, it is not diagnostic of lymphedema.

Case 2

An 18-year-old woman describes a history of larger legs that accelerated during puberty. Despite multiple attempts at diet-induced weight loss and, more recently, medication-induced weight loss, she has been unable to reduce her leg size and has relied on wearing larger boots to conceal her appearance. She now seeks assistance for her enlarged legs, which are limiting her mobility.

On physical examination, the patient's vital signs are within normal limits. Her BMI is 32 kg/m², and she has a gynoid fat distribution with excess skin and dimpling. The Stemmer sign is positive. She has no pain to touch or nodularity in her legs.

Given her likely stage of lipedema, what treatment options should be avoided at this point in her disease course?

- A. Dietary strategies
- B. Complete decongestive therapy
- C. Complete lymphatic therapy
- D. Surgical removal

Answer: D) Surgical removal

This patient likely has class I or II lipedema based on the provided description and is unlikely to have had the disease for more than 10 years, given its onset at puberty. As such, there is ample opportunity to implement a layered approach to prevent progression of lipedema, including dietary strategies (Answer A) and mechanical therapies (eg, complete decongestive therapy [Answer B] and complete lymphatic therapy [Answer C]). Surgical removal (Answer D) of lipedema adipose tissue should be reserved for patients with significant pain or mobility limitations or advanced disease, but guidelines are evolving.

Case 3

A 37-year-old woman with hypertension and type 2 diabetes (hemoglobin A_{1c} = 6.3% [45 mmol/mol]), controlled for 3 years on metformin, presents to the endocrinology clinic

with concern about leg swelling and oozing. She endorses increased leg mass since menarche and, over the past 4 years, has gained significant weight that has been unresponsive to diuretic treatment or GLP-1 receptor agonist therapy. She now reports pain and painful, pearl-like nodules in her legs, along with oozing from her heels.

On physical examination, the patient has a BMI of 41 kg/m² and stable vital signs. Her light-touch sensation remains intact on monofilament testing, and there are mild abrasions on the heel and plantar surfaces with clear fluid oozing. She has excess skin on her thighs and decreased mobility, with difficulty standing from her chair.

Given the patient's presentation with more advanced lipo-lymphedema, what initial therapeutic option should be considered?

- A. Diuretic therapy
- B. Complete decongestive therapy
- C. Another trial of GLP-1 receptor agonist therapy
- D. Surgical removal

Answer: B) Complete decongestive therapy

This patient with lipo-lymphedema would benefit from a layered approach, beginning with mechanical decompression therapies, such as complete decongestive therapy (Answer B), before proceeding to medical or surgical therapies. A vascular medicine evaluation will be important to assess the extent of lymphedema and its effect on lipedema progression. Following vascular evaluation, diuretic therapy (Answer A) may be considered. GLP-1 receptor agonist therapy (Answer C) is generally unable to remove lipedema fat once fibrosis has formed. Surgical removal (Answer D) may be considered after initial mechanical and medical therapies.

Key Learning Points

- Lipedema remains underdiagnosed in the endocrinology community as a disease of loose connective tissue, including adipose tissue,

often overlapping with obesity, characterized by abnormal lower-body fat distribution and pain, and primarily affecting women.

- Lipedema sits at the intersection of endocrinology, given the important hormonal cues to its development, and adipose biology, given the integral role of adipocytes in its development.
- Lipedema does not respond to conventional weight-loss treatments, including GLP-1 receptor agonists or dietary interventions, and treatment often requires a layered approach that includes mechanical, medical, and ultimately surgical interventions.
- Ongoing research aims to identify blood-based and imaging biomarkers that could inform future therapies.

References

1. Ernst AM, Bauer H, Bauer HC, Steiner M, Malfertheiner A, Lipp AT. Lipedema research-quo vadis? *J Pers Med*. 2022;13(1):98. PMID: 36675759
2. Lundanes J, Sandnes F, Gjeilo KH, et al. Effect of a low-carbohydrate diet on pain and quality of life in female patients with lipedema: a randomized controlled trial. *Obesity (Silver Spring)*. 2024;32(6):1071-1082. PMID: 38627016
3. Shavit E, Wollina U, Alavi A. Lipoedema is not lymphoedema: a review of current literature. *Int Wound J*. 2018;15(6):921-928. PMID: 29956468
4. Herbst KL, Kahn LA, Iker E, et al. Standard of care for lipedema in the United States. *Phlebology*. 2021;36(10):779-796. PMID: 34049453
5. Kruppa P, Georgiou I, Biermann N, Prantl L, Klein-Weigel P, Ghods M. Lipedema-pathogenesis, diagnosis, and treatment options. *Dtsch Arztebl Int*. 2020;117(22-23):396-403. PMID: 32762835
6. Tomada I. Lipedema: from women's hormonal changes to nutritional intervention. *Endocrine*. 2025;6(24):1-19.
7. Fife CE, Maus EA, Carter MJ. Lipedema: a frequently misdiagnosed and misunderstood fatty deposition syndrome. *Adv Skin Wound Care*. 2010;23(2):81-94. PMID: 20087075
8. Frank AP, de Souza Santos R, Palmer BF, Clegg DJ. Determinants of body fat distribution in humans may provide insight about obesity-related health risks. *J Lipid Res*. 2019;60(10):1710-1719. PMID: 30097511
9. Felmerer G, Stylianaki A, Hägerling R, et al. Adipose tissue hypertrophy, an aberrant biochemical profile and distinct gene expression in lipedema. *J Surg Res*. 2020;253:294-303. PMID: 32407981
10. Al-Ghadban S, Cromer W, Allen M, et al. Dilated blood and lymphatic microvessels, angiogenesis, increased macrophages, and adipocyte hypertrophy in lipedema thigh skin and fat tissue. *J Obes*. 2019;2019:8747461. PMID: 30949365
11. Wiedner M, Aghajanzadeh D, Richter DF. Differential diagnoses and treatment of lipedema. *Plast Aesthetic Res*. 2020;7(0):10.
12. Mortada H, Alhithlool AW, AlBattal NZ, et al. Lipedema: clinical features, diagnosis, and management. *Arch Plast Surg*. 2025;52(3):185-196. PMID: 40386000

2026 Endocrine Case Management: Meet the Professor is your source for the latest updates in the diagnosis and management of a wide range of endocrine disorders. This valuable resource allows you to evaluate your endocrine knowledge and gain insight into the strategies used by clinical experts.

Features of *2026 Endocrine Case Management: Meet the Professor* include:

- Learning Objectives
- Significance of the Clinical Problem
- Practice Gaps
- Discussion
- Clinical Case Vignettes
- Key Learning Points
- Pertinent Clinical References



Hormone Science to Health

Endocrine Society
2055 L Street NW, Suite 600
Washington, DC 20036 USA
endocrine.org

